

SECURITY SECURITY SECURITY

SECURITY SECURITY SECURITY SECURITY

SECURITY SECURITY SECURITY

SECURITY

Continued from back panel

DIRECTIONS

NOTE: Workers entering treated fields within 24 hours should wear protective clothing

Apply the individual amount of material per acre in sufficient water for good coverage. Repeat applications as necessary but never closer to harvest than specified

BEANS: Mexican Bean Beetle, Leafhoppers, Armyworm, Leaf Roller, Leaf Miner, Red Spiders and Aphids - 2 pounds. Do not apply within 7 days of harvest

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, KALE, MUSTARD GREENS, COLLARDS, TURNIPS: Diamond Back Moth Imported Cabbage Worm, Armyworms, Aphids, Thrips, Vegetable Weevil - 3 pounds. Do not apply within 10 days of harvest except 7 days on Broccoli and Brussels Sprouts

CUCUMBERS, SQUASH AND MELONS: Cucumber Beetle, Melonworm, Pickleworm, Serpentine Leaf Miner, Aphids - 1 to 3 pounds. To control Vine Borer and Stink Bugs apply 3 pounds. Do not apply within 15 days of harvest to Cucumbers and Squash and 7 days to Melon. Do not apply before vining

PEPPER: Aphids and Serpentine Leaf Miner - 2 pounds. Do not apply within 15 days of harvest

ONIONS: Thrips - 2 pounds. Do not apply within 15 days of harvest

POTATO AND TOMATO: Colorado Potato Beetle, Fleabeetle, Serpentine Leaf Miner, Grasshoppers, Whitefly and Aphids - 2 pounds with regular fungicide applications. Do not apply within 5 days of harvest to potatoes and 10 days to tomatoes

Tobacco

Field: Aphids and Suckly - 1 to 2 pounds in sufficient water for coverage. Hornworm - 2 to 3 pounds in sufficient water for coverage. Make no application within 5 days of pruning or 15 days of cutting. Avoid getting wet with plant juices when cutting

Seedbed: Aphids - 1/2 lb. to 25 gallons of water applied at 3 to 5 gallons per 100 square yards. Do not apply within 5 days of transplanting

ORNAMENTALS (Outdoor Only)

Use only on commercially grown field plants. Do not use around homes. Aphids, Whiteflies, Red Spider Mites, Leafhoppers, Thrips, Lace Bug, Bagworm, Cottony Cushion Scale - 1 pound per 100 gallons of water at 150 to 250 gallons per acre. Repeat as necessary. Repeat in 7 to 10 days for Spider Mites and Leafhoppers. Juniper Webworm, Lecanium scale - 1/2 pounds per 100 gallons of water.

Mealybugs, Leaf Tiers, Scale - 1/4 pounds per 100 gallons of water. A repeat application may be necessary in 3 weeks

FRUITS AND NUTS

For control of such orchard pests as Oriental Fruit Moth and Curculio on peaches and Codling Moth on apples, apply sprays at petal fall and follow every 10 to 14 days with additional applications as long as conditions warrant

PEACH (Areas Other than California): Curculio, Oriental Fruit Moth, San Jose Scale (crawler stage only) and Cat Facing Insects - 2 pounds per 100 gallons of water. Fruit Tree Leaf Roller, Cottony

(Continued on other side panel)

STOP! READ THE LABEL

POISON
See back panel for antidote and other precautions



DANGER
Keep out of reach of children

CAN KILL YOU

15%

PARATHION WETTABLE

NOT FOR HOME USE

NET WT. 3 LBS.

Active Ingredient:

Parathion: O, O-diethyl O-p-nitrophenyl phosphorothioate . . . 15.0%

Inert Ingredients: 85.0%

TOTAL 100.0%

EPA REG. NO. 769-77

POISONOUS TO FISH AND WILDLIFE: This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

PROTECT BEES: This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE—DISCLAIMER OF WARRANTY, MERCHANTABILITY AND LIABILITY—If for any reason this product is not used, applied, stored, transported or disposed of in accordance with the instructions set forth on this label, then and thereafter Woolfolk Chemical Works, Ltd., terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness

(Continued from other side panel)

Peach Scale, Green Peach Aphid, and Spider Mites - 1 pound per 100 gallons of water. Peach Tree Borer, Lesser Peach Tree Borer - Use 3 pounds per 100 gallons of water following local recommendations for timing and frequency of applications. Make applications during appearance of the moths in early to late summer. Apply 2 or 3 sprays to trunk from ground to scaffold must timed with moth flight. Do not apply more than 5 pounds of actual parathion per acre

APPLE: Codling Moth, Plum Curculio, Orange Tortrix, San Jose, Pear and Scurfy Scales, Red Banded Leaf Roller (Second and third brood), Woolly Apple Aphid and Mealybug - 1 1/4 pounds per 100 gallons of water. Bud Moth, Red Bug, Fruit Tree Leaf Roller, Raggy Apple Aphid, Green Apple Aphid, Red-Banded Leaf Roller (first brood), Grasshoppers, and European Red Mite, Two-spotted Mite, Pacific Mite, Schonek Mite, Willamette Mite and Clover Mite - 1 pound per 100 gallons of water. For certain mites such as two-spotted and Willamette applications should be repeated at 7 day intervals.

NOTE: Foliage injury to McIntosh, Cortland, Macoun, Kendall, Melba, Golden Delicious and Jonathan apples may result from using rates as listed above

PECAN: Nut Casebearer - Use 2 pounds per 100 gallons of water applied at the time tips of small nuts have turned brown. Black Pecan Aphid, Mites, and Leaf Casebearer - Use 2 pounds per 100 gallons of water - except for mites, use 3 pounds. Repeat applications as necessary. DO NOT APPLY TO PECANS WITHIN 15 DAYS OF HARVEST. DO NOT GRAZE PECAN GROVES WITHIN 15 DAYS OF APPLICATION

MAKE NO APPLICATION TO ABOVE DECIDUOUS FRUIT CROPS WITHIN 14 DAYS OF HARVEST

SMALL FRUITS

GRAPE: Mealybug and Leaf Folder - 1 1/4 pounds per 100 gallons of water. Mites - 1 pound per 100 gallons of water repeated at 5 to 7 day intervals. Make applications before fruit is the size of buckshot or after harvest. Apply 200 gallons of dilute spray per acre. Do not apply within 14 days of harvest

BLUEBERRY: Maggot and Thrips - 1 1/4 pounds per 100 gallons of water and apply at the rate of 175 to 150 gallons per acre. Do not apply within 14 days of harvest

STRAWBERRY: Leaf Roller and Red Spider Mites - 1 to 1 1/4 pounds per 100 gallons of water applied at the rate of 100 to 200 gallons of dilute spray per acre. Repeat at 7 to 10 day intervals. Do not apply within 14 days of harvest

DISPOSAL WARNING

Dispose of waste pesticide and containers by burying in a deep land away from water supplies. Never reuse container

NOTICE

DISCLAIMER OF WARRANTY, MERCHANTABILITY AND LIABILITY

If for any reason this product is not used, applied, stored, transported or disposed of in accordance with the instructions set forth on this label, then and thereafter Woolfolk Chemical Works, Ltd. terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness



CAN KILL IF SWALLOWED

This product can kill you even in small amounts or dust may be fatal

USE ONLY WHEN WEAR PROTECTIVE EQUIPMENT

(1) Wear water-proof or rubber overshoes. Wear mask or respirator of Mines for parathion duty, natural rubber.

WORK

Keep all unprotected parts of body out of treated area or where

Do not rub eyes or mouth in any way, STOP work if you feel ill. Consult a doctor (physician), explain that the victim has been given (see First Aid Treatment) for cannot come, take

IMPORTANT! Before using, wash hands with soap and water. Do not drink or eat

AFTER WORK: Take a shower, using soap and water. Wash work clothes when leaving job. Wash protective equipment with soap and water. Respirator should be cleaned according to instructions

POISON SIGNS

Parathion is a very dangerous poison. It enters the body on contact with skin, clothing, or eyes. Clothing wet with parathion should be removed immediately. Examine for prompt medical treatment

Some of the signs and symptoms of poisoning are: Headache, nausea, vomiting, blurred vision, pin point pupils, etc.

SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY

STOP! READ THE LABEL



DANGER
Keep out of reach of children

CAN KILL YOU

15%

PARATHION
NETTABLE

NOT FOR HOME USE

NET WT. 3 LBS.

0-p-nitrophenyl phosphorothioate	15.0%
	85.0%
TOTAL	100.0%

AND WILDLIFE: This product is toxic to fish and wildlife in treated areas may be killed. Keep out of water. Do not apply when weather conditions favor drift. Do not apply where runoff is likely to occur. Do not clean or leave equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct contact with treated plants. Protective information may be obtained from Agricultural Extension Service.

FORCE OF WARRANTY, MERCHANTABILITY AND FITNESS: Reason this product is not used, applied, stored, transported or disposed of in accordance with the instructions set forth on this label, then and thereafter Woolfolk Chemical Works, Ltd., terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness.

(Continued from other side panel)

PEACH: Scale, Green Peach Aphid, and Spider Mites - 1 pound per 100 gallons of water. Peach Tree Borer, Lesser Peach Tree Borer - Use 3 pounds per 100 gallons of water following local recommendations for timing and frequency of applications. Make applications during appearance of the moths in early to late summer. Apply 2 or 3 sprays to trunk from ground scaffold thus timed with moth flight. Do not apply more than 3 pounds of actual parathion per acre.

APPLE: Codling Moth, Plum Curculio, Orange Tortrix, San Jose, and Scurfy Scales, Red-Banded Leaf Roller (Second and Third Brood), Waxy Apple Aphid and Mealybug - 1 1/2 pounds per 100 gallons of water. Bud Moth, Red Bug, Fruit Tree Leaf Roller, Rosy Apple Aphid, Green Apple Aphid, Red-Banded Leaf Roller (first brood), Grasshoppers, and European Red Mite, Two-spotted Mite, Pacific Mite, Schoone Mite, Willamette Mite and Clover Mite - 1 pound per 100 gallons of water. For certain mites such as two-spotted and Willamette applications should be repeated at 7 day intervals.

NOTE: Foliage injury to McIntosh, Cortland, Macoun, Kendall, Melba, Golden Delicious and Jonathan apples may result from using rates as listed above.

PECAN: Nut Casebearer - Use 2 pounds per 100 gallons of water applied at the time tips of small nuts have turned brown. Black Pecan Aphid, Mites, and Leaf Casebearer - Use 2 pounds per 100 gallons of water - except for mites, use 3 pounds. Repeat applications as necessary. **DO NOT APPLY TO PECANS WITHIN 15 DAYS OF HARVEST. DO NOT GRAZE PECAN GROVES WITHIN 15 DAYS OF APPLICATION.**

MAKE NO APPLICATION TO ABOVE DECIDUOUS FRUIT CROPS WITHIN 14 DAYS OF HARVEST.

SMALL FRUITS

GRAPE: Mealybug and Leaf Folder - 1 1/2 pounds per 100 gallons of water. Mites - 1 pound per 100 gallons of water repeated at 5 to 7 day intervals. Make applications before fruit is the size of buckshot or after harvest. Apply 200 gallons of dilute spray per acre. Do not apply within 14 days of harvest.

BALDFRUIT: Maggot and Thrips - 1 1/2 pounds per 100 gallons of water and apply at the rate of 125 to 150 gallons per acre. Do not apply within 14 days of harvest.

STRAWBERRY: Leaf Roller and Red Spider Mites - 1 to 1 1/2 pounds per 100 gallons of water applied at the rate of 100 to 200 gallons of dilute spray per acre. Repeat at 7 to 10 day intervals. Do not apply within 14 days of harvest.

DISPOSAL WARNING

Dispose of waste pesticide and containers by burying in non-crop land away from water supplies. Never re-use containers.

**NOTICE
DISCLAIMER OF WARRANTY,
MERCHANTABILITY AND LIABILITY**

If for any reason this product is not used, applied, stored, transported or disposed of in accordance with the instructions set forth on this label, then and thereafter Woolfolk Chemical Works, Ltd. terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness.

DANGER—POISON—PRECAUTIONS



CAN KILL YOU IF SWALLOWED
This product can kill you if swallowed even in small amounts; spray mist or dust may be fatal if swallowed.



CAN KILL YOU BY SKIN CONTACT
This product can kill you if touched by hands or spilled or splashed on skin, in eyes or on clothing (liquid goes through clothes).



CAN KILL YOU IF BREATHED
This product can kill you if vapors, spray mist or dust are breathed.

USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE EQUIPMENT AND CLOTHING

- (1) Wear water-proof pants, coat, hat, rubber boots or rubber overshoes.
- (2) Wear safety goggles.
- (3) Wear mask or respirator approved by the U.S. Bureau of Mines for parathion protection.
- (4) Wear heavy duty, natural rubber gloves.

WORK SAFETY RULES

Keep all unprotected persons and children away from treated area or where there is danger of drift. Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. Call a doctor (physician), clinic or hospital—immediately. Explain that the victim has been exposed to parathion and describe his condition. After first aid is given (see First Aid Treatment Section) and if a doctor cannot come, take victim to clinic or hospital.

IMPORTANT! Before removing gloves, wash them with soap and water. Always wash hands, face and arms with soap and water before smoking, eating or drinking.

AFTER WORK, take off all work clothes and shoes. Shower, using soap and water. Wear only clean clothes when leaving job. Do not wear contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Respirator should be cleaned and filter replaced according to instructions included with respirator.

POISON SIGNS (Symptoms)

Parathion is a very dangerous poison. It rapidly enters the body on contact with all skin surfaces and eyes. Clothing wet with this material must be removed immediately. Exposed persons must receive prompt medical treatment or they may die.

Some of the signs and symptoms of poisoning are: Headache, nausea, vomiting, cramps, weakness, blurred vision, pin point pupils, tightness in chest, labored breathing, nervousness, sweating, restlessness.

FIRST AID TREATMENT

Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to parathion and describe his condition.

If breathing has stopped, start artificial respiration immediately and maintain until doctor sees victim. If swallowed and victim is awake (conscious) make him vomit quickly. Induce vomiting by sticking finger down throat or by giving soapy or strong salty water to drink. Repeat until vomit is clear. Never give anything by mouth to an unconscious person. Have victim lie down and keep quiet. See doctor immediately.

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. See doctor immediately.

NOTE TO PHYSICIAN

Antidote—administer atropine sulfate in large doses, TWO to FOUR mg. intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 10 minute intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. **DO NOT GIVE MORPHINE OR TRANQUILIZERS.** Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement. **VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.**

POST TREATED AREA

Consult your State Agricultural Extension Service or Experiment Station regarding posting treated areas.

CONTAINER DISPOSAL

Dispose of waste pesticide and empty containers by burying in non-crop land away from water supplies. Never re-use container.