

ACTIVE INGREDIENT: Dithionon 25.0
INERT INGREDIENTS: 75.0

100, 0-d ethyl 0-p-n trophenyl thiophos. rate
U.S. D. A. Reg. No. 754-93 TOTAL 100.0%



POISON



ANTIDOTE: IF SWALLOWED: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluids clear. Have victim lie down and keep quiet. Call a Physician immediately! Atropine is antidote. IF ON SKIN: In case of contact remove contaminated clothing and immediately wash skin with soap and water.

WARNING: KEEP OUT OF REACH OF CHILDREN.

Poisonous if swallowed, inhaled, or absorbed through skin. Rapidly absorbed through skin. Do not get in eyes or on skin. Wear natural rubber gloves, or protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U.S. Dept. of Agriculture for protection. Keep all unprotected persons out of operating areas until dust where there may be danger of drift. Drifted areas should not be re-entered until drifting has ceased and dust particles have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or drinking. Do not use in or near water.

PROTECT WILDLIFE: Highly toxic to birds, fish, and aquatic life. Do not apply in or near water or adjacent areas. Do not apply in or near water when weather conditions favor drift. Do not feed treated areas to birds.

PROTECT BEES: This product is highly toxic to bees. Do not apply in or near beehives or bees. Protective clothing should be worn. See also U.S. Department of Agriculture, Entomology Service.

DIRECTIONS

FIELD CROPS

COTTON: Aphids, Red Spider Mite, Fleahopper, Garden Webworm - Apply at the rate of 20 to 30 pounds per acre. Repeat as necessary for control. Do not apply within 5 days of hand picking or within 15 days of cotton trash is to be fed to dairy animals or animals being finished for slaughter.

PEANUTS, SOYBEANS: Velvet Bean Caterpillar, Thrips - Apply at the rate of 20 to 30 pounds per acre. Do not apply within 15 days of harvest, cutting or forage use.

TOBACCO: FIELD - Use power dusting equipment - Aphids, Suckly, Hornworm - Apply at the rate of 15 to 30 pounds per acre as necessary for control. Do not apply within 15 days of harvest. Make no application within 5 days of cutting. Avoid getting wet with plant juices when cutting or handling. **GREENHOUSE:** Aphids, Midge, Green Leaf Beetle - Apply at the rate of 10 to 20 pounds per square yard.

VEGETABLES

INTERVAL BETWEEN LAST APPLICATION AND HARVEST

Manufacturers

apply to Potatoes within 5 days of harvest; to Tomatoes and Peas within 10 days of harvest; to Beans, Black-eyed Peas, Rutabaga and Turnips used for roots, Carrots, Celery, Cucumbers, Squash, Melons, Pumpkins, Eggplant, Onions, Peppers, and Radishes within 15 days of harvest; to Cabbage, Broccoli, Brussels Sprouts, Kale, Mustard Greens, Cauliflower, Kohlrabi, Collards, Rutabaga and Turnips used for tops, Endive, Lettuce, Dava, Spinach and Swiss Chard within 21 days of harvest.

BEANS: Mexican Bean Beetle, Fleahopper, Aphid, Red Spider Mite, Leaf Rollers, Armyworms, Leaf Miner - Apply at the rate of 30 to 40 pounds per acre. Repeat at 7-10 day intervals as necessary for control.

BLACK-EYED PEAS: Aphids, Leaf Miner - Apply at 25 to 40 pounds per acre.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, KALE, MUSTARD GREENS, CAULIFLOWER, KOHLRABI, COLLARDS, RUTABAGA, TURNIP: Diamondback Moth, Imported Cabbage Worm, Armyworm, Aphids, Thrips, Vegetable Weevil - Apply at the rate of 30 to 40 pounds per acre. Repeat as necessary for control.

CARROTS: Aphids, Vegetable Weevil - Apply at the rate of 30 to 40 pounds per acre.

CELERY: Armyworms, Aphids - Apply at the rate of 30 to 40 pounds per acre. Do not apply within 10 days of harvest when rates above 30 pounds per acre are used.

CORN: European Corn Borer, Grasshoppers, Armyworm - Apply at the rate of 45 to 55 pounds per acre as necessary for control. Corn Earworm - Apply at the rate of 25 to 35 pounds per acre as they appear and repeat at 3-day intervals as long as insects are present. Repeat immediately after rain. Leaf Apteris, Fleahopper - Apply at the rate of 20 to 40 pounds per acre. Do not apply within 10 days of cutting or forage. Do not feed forage this stage for 15 days after application.

CUCUMBER, SQUASH, MELONS, PUMPKINS: Cucumber Beetle, Fleahopper, Pickeworm, Serpentine Leaf Miner, Aphids, Stink Bug, White Borer - Apply at the rate of 20 to 40 pounds per acre. Repeat as necessary. Do not apply to cucurbits unless plants are well established before plants start to vine. To control Pickeworm and Fleahopper, begin applications with first vines and repeat at 5 to 7 day intervals as necessary.

EGGPLANT: Aphids, Leaf Miner - Apply at the rate of 20 to 30 pounds per acre.

ENDIVE: Aphids, Leafhoppers - Apply at the rate of 35 to 40 pounds per acre.

LETTUCE: Aphids, Leafhoppers - Apply at the rate of 35 to 40 pounds per acre.

OKRA: Aphids - Apply at the rate of 20 to 30 pounds per acre.

ONIONS: Thrips - Apply at the rate of 25 to 35 pounds per acre. Repeat as necessary.

PEAS: Aphids, Pea Moth - Apply at the rate of 25 to 35 pounds per acre. Do not feed forage or silage for 15 days after last application.

PEPPERS: Aphids, Serpentine Leaf Miner - Apply at the rate of 20 to 30 pounds per acre.

RADISHES: Aphids, Leaf Miners - Apply at the rate of 30 pounds per acre.

SPINACH, SWISS CHARD: Aphids - Apply at the rate of 20 to 30 pounds per acre.

WOOLFOLK CHEMICAL WORKS, LTD.

POISONOUS TO HUMANS AND DOMESTIC ANIMALS
POTATOES, TOMATOES: Colorado Potato Beetle, Flea Beetle, Fleahopper, Serpentine Leaf Miner, Aphids, Grasshopper, Leaf Miner
Apply at the rate of 25 to 35 pounds per acre.

FORAGE CROPS

VETCH, ALFALFA, CLOVER: Aphids, Armyworms, Grasshoppers, Blister Beetles, Alfalfa Caterpillar - Apply at the rate of 25 to 40 pounds per acre. Do not apply within 15 days of harvest or pasturing.

WHEAT, OATS, BARLEY: Aphids, Armyworms, Grasshoppers, Blister Beetle - Apply at the rate of 25 to 40 pounds per acre. Do not apply within 15 days of harvest or pasturing.

ORNAMENTALS

APHIDS, WHITEFLIES, RED SPIDER MITES: Apply at the rate of 20 to 35 pounds per acre. Repeat as necessary. Repeat in 7 to 10 days for control of red spider mites.

MEALYBUGS, LEAF TIERS: Apply at the rate of 30 to 40 pounds per acre. Repeat application in 3 weeks if necessary. Apply this material outdoors only.

CAUTION: This product should be used only by properly equipped personnel on commercial plantings, never on home plantings.

FRUIT AND NUTS

BLACKBERRIES: Red-necked Cane Borer - Apply at the rate of 30 to 35 pounds per acre before bloom. Do not apply after edible parts start to form.

BLUEBERRIES: Maggot, Thrips - Apply at the rate of 30 pounds per acre. Apply only before fruit sets or after harvest.

BOYSENBERRIES, NEWBERRIES: Red Spider Mites, Aphids - Apply at the rate of 20 to 35 pounds per acre. Repeat at 7 to 10 day intervals as necessary. Apply only before bloom or after harvest.

GRAPES: Red Spider Mite, Leaf Folder, Mealybug - Apply at 40 pounds per acre. Repeat in 5 to 7 days for red spider. Do not apply within 14 days of harvest.

STRAWBERRIES: Red Spider Mite, Aphids - Apply at the rate of 20 to 35 pounds per acre. Repeat in 7 to 10 days for red spider. Do not apply within 14 days of harvest.

PECAN: Black Pecan Aphid - Apply to thoroughly cover the trees. Make no application after shucks have started to open. Do not feed vegetation under nut trees treated with this material to dairy or beef animals.

DISPOSAL WARNING: Dispose of waste pesticide and containers by burying in non-crop land away from water supplies. Never re-use container.

NOTICE DISCLAIMER OF WARRANTY, MERCHANTABILITY AND LIABILITY

If for any reason this product is not used, applied, stored, transported or disposed of in accordance with the instructions set forth on this label then and thereafter Woolfolk Chemical Works, Ltd. terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness. The failure to use, apply, store, transport or dispose of this product in accordance with the instructions set forth on the label shall then and thereafter terminate and end Woolfolk Chemical Works, Ltd.'s liability for injury to crops, animals, personal property and real property of every kind and description, and for injury to any person or persons, and for any consequential damages arising out of liability to the foregoing set forth. There shall be no warranties, whether written or oral, made by Woolfolk Chemical Works, Ltd. in connection with this product.

Fort Valley, Ga.