SECURITY

ZINEB DUST

FOR TOBACCO BLUE MOLD, WATERMELON ANTHRACNOSE, TOMATO BLIGHT, OTHER DISEASES

ACTIVE INGREDIENT: Zineb Zinc ethylenebisdithiocarbamate ..... [illegible]
INERT INGREDIENTS: ..... [illegible]

U. S. D. A. Reg. No. 769-106 TOTAL 1[illegible]

CAUTION: KEEP OUT OF REACH OF CHILDREN. May cause irritation of nose, throat, eyes, and skin. Do not breathe dust. In case of skin contact wash with water; for eyes, get medical attention.

DIRECTIONS

PLANT BEDS

TOBACCO: Blue Mold - Dust small plants at the rate of 1½ to 3 pounds per 100 square yards of plant bed. Increase the rate to 3½ pounds when plants are large. At first sign of blue mold dust at the maximum rate regardless of size of plants. Begin dusting as soon as blue mold is first reported in the area or when plants are the size of a dime. Continue until the plants are set in the field or blue mold has run its course. Dust twice weekly and after every rain even if 3 or 4 dusts per week are required.

CABBAGE, BROCCOLI, CAULIFLOWER: Downy Mildew - Apply at the rate of 1½ to 3 pounds per 100 square yards of bed at weekly intervals.

VEGETABLES

APPLICATION: Apply at the rate of 20 to 35 pounds of dust per acre depending upon the size of the crop. Begin application at first sign of or prior to appearance of disease. Repeat at 1 to 2-week intervals as necessary for control unless otherwise noted.

DAYS BETWEEN LAST APPLICATION AND HARVEST: [illegible] beans, beets, broccoli, Brussels sprouts, cabbage, cauliflower, collards, [illegible] mustard greens, spinach, Swiss chard or turnips within [illegible] days of harvest. [illegible] can be made to cantaloupes, cucumbers, eggplant, [illegible] tomatoes and watermelons up to the day of harvest.

BEANS: Downy Mildew, Rust, Anthracnose - Begin when first [illegible] appear.

BEETS, KALE, MUSTARD GREENS, SWISS CHARD, TURNIPS: Cercospora leaf spot, Downy Mildew.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, COLLARDS: Downy Mildew, Alternaria leaf blight.

CANTALOUPES, CUCUMBERS, MUSKMELONS, PUMPKINS, SQUASH, WATERMELONS: Downy Mildew, Alternaria blight, Anthracnose, Gummy stem blight.

LETTUCE, SPINACH: Downy Mildew [illegible] weekly intervals.

EGGPLANT: Phytophthora, Anthracnose, Ripe Rot - Apply at 10-day intervals when disease is evident. Repeat as necessary throughout the season.

TOMATOES: Early and Late blight, Anthracnose, Stemphylium and Septoria leaf spots, Cladosporium gray mold - Begin applications when disease is reported in area, and repeat applications at 7 to 10-day intervals or oftener if blight conditions are severe.

FRUITS

GRAPES: Downy Mildew, Black rot, Ripe rot - Apply at the rate of 25 to 30 pounds per acre. Apply just before bloom, just after bloom, 10 days after bloom, and 2 to 3 weeks later, or follow local spray schedule. Do not apply within 7 days of harvest.

ORNAMENTALS

AZALEAS: Petal blight - Dust at the rate of 20 - 25 pounds per acre and apply 2 to 3 times a week while flowers are opening. Direct dust into flowers and on ground beneath the plants.

GLADIOLUS: Botrytis blight, Curvularia, Stemphylium - Apply at the rate of 20 to 25 pounds per acre at weekly intervals, or oftener under severe disease conditions, from early stages until spikes begin to open.

SNAPDRAGONS, CHRYSANTHEMUMS, CARNATIONS, HOLLYHOCKS: Rust, Anthracnose - Apply at the rate of 20 to 25 pounds per acre. Begin when disease conditions threaten and apply at weekly to 10-day intervals. For control of Botrytis petal spot and Ascochyta may make applications twice weekly during the [illegible] season.

ROSES: Black spot - Apply 20 to 25 pounds per acre when disease first [illegible] and repeat at weekly intervals.

DISPOSAL WARNING

[illegible]

NOTICE
DISCLAIMER OF WARRANTY, MERCHANTABILITY AND LIABILITY

[illegible] Chemical Works, Ltd. terminates, ends and excludes all warranties [illegible]

Manufacturers **WOOLFOLK CHEMICAL WORKS LTD.** Fort Valley, Ga.