

SECURITY

769-136
AUG 16 1974
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER

TOX-SOL-6

Comment

CONTAINS 6 POUNDS TOXAPHENE PER GALLON

ACTIVE INGREDIENTS:

*Toxaphene	59.0%
Petroleum Hydrocarbons	31.8%
Xylene	4.8%
INERT INGREDIENTS:	4.4%

E. P. A. Reg. No. 769-136	TOTAL	100.0%
*Technical Chlorinated Camphene (Chlorine content 67% to 69%)		

WARNING! KEEP OUT OF REACH OF CHILDREN. May Be Fatal If Swallowed!

Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after using. Do not store near feed or food products. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

DIRECTIONS

PROTECT WILDLIFE: This product is poisonous to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on the label. Birds feeding on treated areas may be killed.

PROTECT BEES: This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not feed treated crop or crop refuse to dairy animals or animals being finished for slaughter, or to poultry. Do not allow grazing of treated crops by dairy animals or animals being finished for slaughter. Do not sell or ship forage treated with this material.

PECANS

To control pecan weevil use at the rate of 2 gallons to 500 gallons of water (3.2 pints per 100). Do not apply more than 400 gallons per acre. Follow locally issued spray schedule for number and frequency of applications. Usually two applications are necessary, the first about July 30 in the latitude of Fort Valley, Ga., or when weevils begin to appear as determined by jarring, and a second application should follow in 10 days to 2 weeks. Do not apply after shucks begin to open. Do not graze livestock in treated groves.

LIVESTOCK

Do not use on dairy animals. Do not use in excess of recommended rate. Concentrations of twice the recommended rate may be fatal to young animals. Do

not apply to animals within 28 days of slaughter. Do not use this product in dips.

BEEF CATTLE: Horn flies, Lice and Ticks - Use 5-1/3 pints per 100 gallons of water (1/2 cup to 5 gallons of water) and spray at the rate of 2 quarts per animal. Apply at 3 week intervals or as needed for horn flies and lice.

SHEEP AND GOATS: Lice, Keds (Sheep tick) - Use 5-1/3 pints per 100 gallons of water (1/2 cup to 5 gallons of water) and spray thoroughly. Repeat treatment in 2 to 3 weeks if necessary.

VEGETABLES

Do not apply to or allow drift to reach cucumbers, squash, or melons or injury may result.

COWPEAS: Cowpea curculio - Apply at the rate of 4 pints per acre in sufficient water for coverage. Make the first application at the time of bloom or when first small peas appear. Make 2 to 4 applications at 4 to 5 day intervals. Do not apply to peas to be used as green snaps after pods begin to form. Do not feed treated units to dairy animals or animals being finished for slaughter.

TOMATOES, PEPPERS, EGGPLANT: Hornworms, Blister beetles, and Pepper weevils on peppers - Apply at the rate of 2-2/3 to 4 pints per acre on peppers and eggplant and 2-2/3 to 5-1/3 pints on tomatoes in sufficient water for coverage. Do not apply to tomatoes within 7 days of harvest or to peppers and eggplant within 5 days of harvest. Remove all residue at harvest by washing or other effective means.

CUTWORMS: On beans, broccoli, cabbage, cauliflower, celery, eggplant, kale, peppers, and tomatoes. Apply 2-2/3 pints per acre in sufficient water for coverage to soil surface before planting or to soil surface immediately after setting plants. Do not apply when plants are past seedling stage.

COTTON

THRIPS: For early season control apply at the rate of 1 to 2 pints per acre in sufficient water for coverage, usually 2 to 3 gallons with one nozzle per row. Make the first application as soon as cotton is up to a stand and a second application 7 days later.

SOYBEANS AND PEANUTS

CUTWORMS, FALL ARMYWORM, VELVETBEAN CATERPILLAR: Apply at the rate of 2-2/3 pints per acre in sufficient water for coverage. Make application as soon as small worms are found. Do not apply to soybeans within 21 days of harvest.

BARLEY, OATS, WHEAT, RYE, CORN, SORGHUM, ARMYWORM, FALL ARMYWORM AND GRASSHOPPERS: Apply at the rate of 2 to 2-2/3 pints per acre in sufficient water for coverage. Do not apply to sorghum within 28 days of harvest. Do not apply to sorghum more than once after heads begin to form. Do not ensile treated forage.

FLOWERS, LAWNS AND ORNAMENTALS

TICKS AND CHIGGERS: Apply at the rate of 4 to 5-1/3 pints per acre (1/2 cup to 2000 square feet) in sufficient water for coverage. Treat infested areas, paying particular attention to weedy areas. Repeat in 4 weeks or as necessary for control.

FALL ARMYWORM AND GRASSWORM: On lawns, apply at the rate of 5-1/3 pints per acre (1/2 cup to 2000 square feet) using sufficient water for coverage.

FOLIAGE FEEDING INSECTS: Bagworm on arbovitae, juniper, soruce, hemlock, cedar and pyracantha; Blister beetles on aster, clematis, chrysanthemum, mimosa and zinnia; Cankerworm on birch, elm and oak; Catalpa worm; Elm sawfly; Elm beetles on oak, pecan and rose; Mimosa webworm; Fall Webworm and Tea Caterpillar on beech, birch, oak, petal, poplar, rose and willow; Sawfly on pine, soruce and honeysuckle - Use 3 to 4 pints per 100 gallons of water (2 tablespoons per gallon) and apply for thorough coverage. Repeat as necessary.

CUTWORMS: Apply to the surface of flower beds before planting, at the rate of 1/2 cup to 2000 square feet using sufficient water for good coverage.

DISPOSAL WARNING

Dispose of waste pesticide and spillage by burying in non-crop land away from water supplies. Containers should be perforated or crushed and buried with waste.

NOTICE

DISCLAIMER OF WARRANTY, MERCHANTABILITY AND LIABILITY

If for any reason this product is not used, applied, stored, transported or disposed of in accordance with the instructions set forth on this label, then and thereafter Woolfolk Chemical Works, Inc. terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness.

WOOLFOLK CHEMICAL WORKS, INC.

MANUFACTURERS 370405 FORT VALLEY, GEORGIA

ACCEPTED WITH COMMENTS

NET WEIGHT



FIVE POUNDS

TOMATO DUST

ACTIVE INGREDIENTS:

Trivalent Arsenate, $Ca_3As_2O_7$ 6.0%
Copper, in the form of Tri-basic Copper Sulphate, as metallic 2.0%

INERT INGREDIENTS: 92.0%

U. S. D. A. Reg. No. 769-149 TOTAL 100.0%

Total arsenic as metallic, 0.25%. Water soluble arsenic as metallic, not more than 0.49%.

DANGER: KEEP OUT OF REACH OF CHILDREN. Poisonous if Swallowed. Avoid contact with skin, eyes, or clothing. Wash thoroughly after using. Avoid breathing dust. Keep away from pets, livestock, feed and foodstuffs.

PROTECT WILDLIFE: This product is toxic to wildlife. Birds and other wildlife in treated areas may be harmed. Apply this product only as specified on this label.



POISON



ANTIDOTE: Calcium Plus, 1/2 tsp. immediately. Give a teaspoonful of salt in a glass of warm water and repeat until vomit turns clear. Then give two tablespoons of Epsom Salts in Milk of Magnesia in water, and plenty of milk and water. Have victim lie down and keep quiet.

DIRECTIONS

TOMATOES: Tomatoes: Early Blight, Late Blight, Navel Rot, Septoria Leaf Spot, Leaf Mold, Gray Leaf Spot, Early Blight, Potato Beetle, Tomato Fruitworm, Hornworm, Tomato Pinworm. Use a good hand sprayer for application. For spray applications purchase the 5 lb. Security Brand Duster container and use it as directed on the label as necessary. Start dusting as soon as plants become well established. Best results are obtained when dusting is done when the air is calm and when dew is on the plants. Best results are obtained when dusting is done in the evening and in the morning. Direct the dust to the undersides of leaves. To control the fruit fly attack by the tomato fruitworm, start dusting intervals beginning with the first fruit. The extent of dusting will depend on the extent of the pest. The amount of application will depend on the type of the plants. The amount of dusting should be made on the application of the dust to the leaves and the fruit.

CAUTION: Do not use on sensitive plants.

DISPOSAL WARNING: Do not use on sensitive plants.

NOTICE

DISCLAIMER OF WARRANTY, MERCHANTABILITY AND LIABILITY

This product is sold as is, without warranty, and the user assumes all liability for its use. The user agrees to indemnify and hold the manufacturer harmless from all claims, damages, and expenses, including reasonable attorney's fees, arising out of the use of this product, whether or not such claims, damages, and expenses are caused in whole or in part by the negligence of the manufacturer.

WOOLFOLK CHEMICAL WORKS, LTD.
Manufacturers Fort Valley, Ga.

May 19, 1969
769-149
769-149