

ORNAMENTALS

APHIDS, THIRPS, LEAFMINERS, CUCUMBER BEETLES: On Boxwood, Azalea, Rose, Chrysanthemum, Larkspur, and other flowers, shrubs, and trees. Use one pint per 100 gallons of water or 1 teaspoonful to 1 gallon of water.

LACEBUG: On Azalea, Pyracantha, and Rhododendron, use 1½ pints per 100 gallons of water or 1½ teaspoons to 1 gallon. Apply in spring when immature insects first appear and repeat as necessary.

PINE BARK BEETLES, TURPENTINE BEETLE: Use 1 pint in 5 gallons of water and apply to trunk of pines to a height of 6 feet at the rate of 1 gallon to 50 square feet of bark surface. Apply at first sign of attack and repeat in 3 to 6 months if necessary.

DOGWOOD BORER: Use 3 pints per 100 gallons of water or 1 tablespoon to 1 gallon and apply to trunk and lower limbs in May and June before borers enter the trunk. Apply thoroughly, particularly to injured areas and old borer wounds. Repeat application 2 or 3 times at 10 day intervals.

LAWNS

TICKS, CHIGGERS, FLEAS, LEAFHOPPERS: Use 1 pint to 10,000 square feet or 3 tablespoons to 1000 square feet in sufficient water for good coverage. Repeat application if necessary. Birds feeding on treated areas may be harmed.

DISPOSAL WARNING

Dispose of waste insecticide and spillage by burying in non-crop land away from water supplies. Perforate or break empty container and bury with waste.

NOTICE

The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

Net Contents

SECURITY

1 Gallon

20% LINDANE EMULSIFIABLE CONCENTRATE

CONTAINS 1.65 POUNDS LINDANE PER GALLON

ACTIVE INGREDIENTS:

Gamma Isomer of Benzene Hexachloride from Lindane 20.0%
Aromatic petroleum derivative solvent 65.6%

INERT INGREDIENTS:

U. S. D. A. Reg. No. 769-159 TOTAL 100.0%

CAUTION! KEEP OUT OF REACH OF CHILDREN.

Harmful if Swallowed. May be absorbed through skin. Avoid inhalation and skin contact. In case of contact, wash immediately with soap and water. Avoid contamination of feed and food-stuffs. Do not use on household pets or humans.

PROTECT WILDLIFE: Toxic to fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on the label. Do not use pour, spill or store near heat or open flame.

PROTECT BEES: This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Woolfolk Chemical Works, Ltd.

Manufacturers

Fort Valley, Ga.

DIRECTIONS LIVESTOCK

CAUTION: Do not use on dairy animals. Do not use on emaciated, sick, overheated or lactating animals. Do not treat animals in cold stormy weather. Do not spray animals within 30 days or dip animals within 60 days of slaughter. Thoroughly stir dipping vats before using. If any separation of the emulsion occurs, do not use the dip.

BEEF CATTLE: LICE AND TICKS: Use 1½ pints per 100 gallons of water. Thoroughly spray entire animal. Repeat treatment in 2 to 3 weeks if necessary. **SARCOPTIC MANGE AND SCAB MITES:** Spray using 2 pints per 100 gallons of water. Make only 2 applications, 7 days apart. Do not apply to calves under 6 months of age at this dosage.

SWINE: LICE AND SARCOPTIC MANGE MITES: Use 2 pints per 100 gallons of water and apply as a spray or dip. For mange mites, repeat treatment in 2 weeks if necessary. Do not treat sows within two weeks of farrowing nor for at least three weeks thereafter.

SHEEP and GOATS: TICKS, KEDS, LICE: Use 2 pints per 100 gallons of water as a spray or use 1 pint per 100 gallons in a dip. Repeat treatment 2 to 3 weeks if necessary.

RESIDUAL SPRAYS

STABLE FLY, MOSQUITOES, ROACHES, ANTS: Inside and outside of barns, hog pens, stables and other buildings. Use 1½ to 2 pints to 10 gallons of water and apply at the rate of 1 gallon to 1000 square feet of surface. Thoroughly cover roosting surfaces for flies and mosquitoes. Apply to cracks, crevices, and other hiding places for control of roaches and ants. Repeat applications as necessary. Do not apply in milkrooms or poultry houses. Do not contaminate animal feed or water.

TICKS and CHIGGERS: Use 2 quarts per 100 gallons of water and spray weeds and underbrush harboring nests. Repeat applications as necessary. Do not use on food crops or grazing areas.

ACCEPTED WITH COMMENTS