


5%

Malathion Dust

NET WT. 4 LBS.




THIS PACKAGE CONTAINS:

ACTIVE INGREDIENT	
*Malathion	5.0%
INERT INGREDIENTS	95.0%
Total	100.0%


EPA Reg. No. 769-161

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

**CONTROLS
MEXICAN
BEAN BEETLE**



ADULT



LARVA





CAUTION: Irritant if swallowed. Avoid breathing of dust. Avoid contact with skin. Wash thoroughly. Avoid contamination of feed and foodstuffs.

PROTECT WILDLIFE

PROTECT BEES: This product is highly toxic to bees exposed to direct treatment or residues on blossoms. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTE TO PHYSICIANS: Malathion may cause cholinesterase inhibition. Atropine is antidote.

DIRECTIONS

To control the insects listed below dust at the rate of 25 to 30 pounds per acre or 2 ounces to each 50 feet of row in the garden. Repeat the applications as necessary. Usually two or more applications at 5 to 7 day intervals will be necessary. To control Mexican Bean Beetle direct the dust to the underside of the leaves.

VEGETABLES

BEANS: Aphids, Leafhoppers, Mexican Bean Beetle, Spider Mites.

CABBAGE, BROCCOLI, TURNIP and MUSTARD: Aphids, Imported Cabbage Worm.

CUCUMBERS, SQUASH, and MELONS: Aphids and Spider Mites. Do not apply to plants unless they are dry or injury may result.

EGGPLANT: Spider Mites and Eggplant Lace Bug.

ONION: Thrips.

PEAS: Aphids and Pea Weevil.

PEPPERS: Aphids.

POTATO and TOMATO: Aphids, Spider Mites and Leafhoppers.

Do not use on Peas, Cucumbers, Melons, or Squash within 1 day of harvest. Broccoli, Eggplant, Onions, Peas, Peppers, Tomatoes, Turnips or Potatoes, within 3 days of harvest. Onion Cabbage or Mustard within 7 days of harvest.

BEEF AND DAIRY CATTLE

HORNFLIES: Apply 1/2 oz. (3 teaspoons) of 5% dust to the back and neck of each animal. Repeat at 10-14 day intervals. Do not apply to dairy cattle later than 5 days before milking. Do not contaminate water troughs.

POULTRY

CHICKENS, DUCKS, GESE and TURKEYS: To control Northern Fowl Mite, use dust as a supplement to premix treatment for Chicken Red Mite. Dust and dust on birds by hand. Do not shake or stir dust. Repeat applications in 4 to 8 weeks or when necessary.

PREMISE TREATMENT: To control Northern Fowl Mite, Chicken Red Mite, and House Fly apply at the rate of 1 lb. to 10-60 sq. ft. of floor and finish space and to nests, roosts and other areas. Repeat application as necessary.

AVOID CONTAMINATION OF FEED AND FEEDING TROUGHS AND WATER FOUNTAINS

DOGS AND CATS

FLEAS: Apply dust enough dust to the body of the animal. Apply to the back and neck to the head with the hand. Wash hands thoroughly. Repeat applications as necessary for control. Do not use more often than once weekly. Do not treat animals under 4 weeks of age.

FRUITS

BLACKBERRY and LOGAN BERRY: To control Leafhopper, apply dust to the leaves. Do not apply within 1 day of harvest. Apply 1/2 oz. (3 teaspoons) of 5% dust per 50 feet of row. To control Aphids, apply dust at the rate of 25 to 30 pounds per acre or 2 ounces to each 50 feet of row. Do not apply within 1 day of harvest.

APRICOTS: Repeat applications for control. Do not apply within 7 days of harvest.

ORNAMENTALS

SPIDER MITES and EGGPLANT LACE BUG: Apply dust to ornamental plants thoroughly. Repeat applications as necessary for control.

SPIDER MITES: Apply dust to certain ferns, including Boston, Maiden Hair and Adiantum, and to plants of CRASSULA.


SPIDER MITES: Apply dust to certain ferns, including Boston, Maiden Hair and Adiantum, and to plants of CRASSULA.

WASTE DISPOSAL: Do not use of waste pesticide. Do not store in a cool and safe place. Do not use in a cool and safe place.


WINDFOLK CHEMICAL WORKS, INC.

Fort Valley, Georgia 31030

CONTROLS MEXICAN BEAN BEETLE



ADULT



LARVA