

Under the Federal Insecticide, Fungicide, and Rodenticide Act.
MAY 17 1976
ACCEPTED

SECURITY

10% CHLORDANE GRANULATED KILLS FIRE ANTS

ACTIVE INGREDIENT:			
*Technical Chlordane	10.0%	
INERT INGREDIENTS:			
.....		90.0%	
EPA Reg. No. 769-177	EPA Est. 769-GA-1	TOTAL	100.0%

*Equivalent to 6% Octachloro-4, 7-methanotetrahydroindane and 4% related compounds.

KEEP OUT OF REACH OF CHILDREN.

CAUTION!

SEE ADDITIONAL PRECAUTIONARY STATEMENTS BELOW.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

KEEP OUT OF REACH OF CHILDREN. Avoid inhaling dust. Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid contamination of feed and foodstuffs. If spilled on skin, wash thoroughly with soap and water. Do not allow livestock to graze treated areas. Do not feed treated forage to dairy animals or animals being finished for slaughter. Keep children and pets off treated areas until this material is washed into the soil and the grass is dry.

HAZARD TO FISH AND WILDLIFE:
This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of ponds, lakes, or streams. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FIRE ANTS

To control fire ants in noncropland only (lawns, ornamental turf, cemeteries, golf courses, parks, industrial sites, roadsides and similar noncropland).

FOR MOUND TREATMENT ONLY - Apply 1-2/3 to 3-1/3 cups Chlordane 10% (equivalent to 0.07 to 0.14 lbs. technical chlordane) to entrance of nest as well as to an area 3 feet around mound.

Treated areas should be examined 3 weeks after treatment and surviving colonies retreated. Do not graze livestock on treated areas within 1 year of treatment application. Do not contaminate feed or food crops.

WHITE-FRINGED BEETLE LARVAE

(Limited to the 8 Southeastern states of Alabama, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina, and Tennessee.)

BROADCAST TREATMENT - Cropland to be planted to beans, cabbage, potatoes, sweet potatoes, and turnips can be treated prior to planting. Apply to soil surface at the rate of 2 1/4 lbs. per 1000 square feet and work into the upper 2 to 3 inches of soil.

JAPANESE BEETLE LARVAE

(Limited to use under Federal/State Quarantine Programs.)

Turf and Nursery Stock - Apply to soil surface at the rate of 2 1/4 lbs. per 1000 square feet.

STORAGE - Do not store near feed or foodstuffs. Keep out of reach of children.

DISPOSAL - Dispose of waste pesticides and container in a safe place away from water supplies. Never re-use container.

WARRANTY STATEMENT

If for any reason this product is not used, applied, stored or disposed of in accordance with instructions set forth on this label, then and thereafter Woolfolk Chemical Works, Inc. terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness.

Manufacturers

WOOLFOLK CHEMICAL WORKS, INC.

Fort Valley, Ga.