

NET WEIGHT



50 POUNDS

2%

ENDRIN DUST

769-179

ACCEPTED

MAY 3 1973

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR THE CONTROL OF REGISTERED UNDER NO. 769-179 SUBJECT TO ATTACHED COMMENTS.

ACTIVE INGREDIENT:

Endrin (Hexachloroepoxyoctahydro-endo, endo-dimethanonaphthalene) 2.0%

INERT INGREDIENTS:

98.0%

EPA Reg. No. 769-179

TOTAL 100.0%

WARNING! KEEP OUT OF REACH OF CHILDREN.

Hazardous by Swallowing, Inhalation or Skin Contact! Rapidly absorbed through skin! Do not get in eyes, on skin, or on clothing. Do not breathe dust. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking; wear clean clothing. Wear clean synthetic rubber gloves and a mask or respirator of a type passed by the U. S. Bureau of Mines for endrin protection.

CAUTION: Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Workers entering fields within five days after treatment should be protected. Keep away from domestic animals. Do not contaminate feed and foodstuffs. Do not graze dairy animals or animals being finished for slaughter on treated cotton fields.

PROTECT WILDLIFE: This product is poisonous to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on the label. Birds feeding on treated areas may be killed.

PROTECT BEES: This product is toxic to bees and should not be applied when bees are actively visiting the area.

DIRECTIONS

COTTON: Boll Weevil, Cotton Bollworm, Cotton Leafworms, Thrips, Cutworms, Fleahoppers, Lygus Bugs and Grasshoppers: To control cutworms apply at the rate of 10 to 20 pounds per acre. To control Thrips apply 5 to 7 pounds per acre at 7-day intervals for three applications beginning when first two true leaves spread. For early season control of overwintered boll weevils, fleahoppers, tarnished plant bug and rapid plant bug, apply

at the rate of 10 pounds per acre making 3 to 4 applications at 7-day intervals beginning at first squaring.

For mid-season control, apply at 10 to 15 pounds per acre. Repeat applications at 5-day intervals until infestation is brought under control. Usually 4 applications will be necessary. Repeat a series of 4 applications at 5-day intervals if boll weevils reappear in sufficient numbers to cause damage. For late season control of migrating boll weevils, apply at the rate of 10 to 25 pounds per acre at 4-day intervals.

Anytime bollworms are a threat, when as many as 4 to 5 small worms and eggs are found per 100 terminals examined, apply 15 pounds of dust per acre. Direct the dust downward into the tops of the plants where young worms are hatching. Repeat applications at 5-day intervals until infestation is controlled.

DISPOSAL WARNING

Dispose of waste pesticide and spillage by burying in non-crop land away from water supplies. Containers should be perforated or crushed and buried with waste.

NOTICE

DISCLAIMER OF WARRANTY, MERCHANTABILITY AND LIABILITY

If for any reason this product is not used, applied, stored, transported or disposed of in accordance with the instructions set forth on this label, then and thereafter Woolfolk Chemical Works, Inc. terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness. The failure to use, apply, store, transport or dispose of this product, in accordance with the instructions set forth on the label, shall then and thereafter terminate and end Woolfolk Chemical Works, Inc.'s liability for injury to crops, animals, personal property and real property, of every kind and description, and for injury to any person or persons, and for any consequential damages arising out of injury to the foregoing set forth. There are no warranties which extend beyond the description on this label.

Manufacturers

WOOLFOLK CHEMICAL WORKS, INC.

Fort Valley, Ga.

NET WEIGHT



50 POUNDS

2%

ENDRIN DUST

769-179

ACCEPTED

MAY 3 1973

U.S. DEPARTMENT OF AGRICULTURE
FUNGICIDE AND PESTICIDE ACT
FOR THE REGISTERED
EPA REG. NO. 769-179 SUBJECT
TO ATTACHED COMMENTS.

ACTIVE INGREDIENT:

Endrin (Hexachloroepoxyoctahydro-endo, endo-dimethanonaphthalene) 2.0%

INERT INGREDIENTS: 98.0%

EPA Reg. No. 769-179

TOTAL 100.0%

WARNING! KEEP OUT OF REACH OF CHILDREN.
Hazardous by Swallowing, Inhalation or Skin Contact!
Rapidly absorbed through skin. Do not get in eyes, on skin, or on clothing. Do not breathe dust. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking; wear clean clothing. Wear clean synthetic rubber gloves and a mask or respirator of a type passed by the U. S. Bureau of Mines for endrin protection.

CAUTION: Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Workers entering fields within five days after treatment should be protected. Keep away from domestic animals. Do not contaminate feed and foodstuffs. Do not graze dairy animals or animals being finished for slaughter on treated cotton fields.

PROTECT WILDLIFE: This product is poisonous to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on the label. Birds feeding on treated areas may be killed.

PROTECT BEES: This product is toxic to bees and should not be applied when bees are actively visiting the crop.

DIRECTIONS

COTTON: Boll Weevil, Cotton Bollworm, Cotton Leafworms, Thrips, Cutworms, Fleahoppers, Lygus Bugs and Grasshoppers: To control cutworms apply at the rate of 10 to 20 pounds per acre. To control Thrips apply 5 to 7 pounds per acre at 7-day intervals for three applications beginning when first two true leaves spread. For early season control of overwintered boll weevils, fleahoppers, tarnished plant bug and rapid plant bug, apply

at the rate of 10 pounds per acre making 3 to 4 applications at 7-day intervals beginning at first squaring.

For mid-season control, apply at 10 to 15 pounds per acre. Repeat applications at 5-day intervals until infestation is brought under control. Usually 4 applications will be necessary. Repeat a series of 4 applications at 5-day intervals if boll weevils reach in sufficient numbers to cause damage. For late season control of migrating boll weevils, apply at the rate of 10 to 25 pounds per acre at 4-day intervals.

Anytime bollworms are a threat, when as many as 4 to 5 small worms and eggs are found per 100 terminals examined, apply 15 pounds of dust per acre. Direct the dust downward into the tops of the plants where young worms are hatching. Repeat applications at 5-day intervals until infestation is controlled.

DISPOSAL WARNING

Dispose of waste pesticide and spillage by burying in non-crop land away from water supplies. Containers should be punctured or crushed and buried with waste.

NOTICE

DISCLAIMER OF WARRANTY, MERCHANTABILITY AND LIABILITY

If for any reason this product is not used, applied, stored, transported or distributed in accordance with the instructions set forth on this label, then and thereafter Woolfolk Chemical Works, Inc. terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness. The failure to use, apply, store, transport or dispose of this product, in accordance with the instructions set forth on the label, shall then and thereafter terminate and end Woolfolk Chemical Works, Inc.'s liability for injury to crops, animals, personal property and real property, of every kind and description, and for injury to any person or persons, and for any consequential damages arising out of injury to the foregoing set forth. There are no warranties which extend beyond the description on this label.

Manufacturers

WOOLFOLK CHEMICAL WORKS, INC.

Fort Valley, Ga.