

ACCEPTED  
29 APR 1980

BEST DOCUMENT AVAILABLE  
SECURITY

# WETTABLE SULPHUR

## ACTIVE INGREDIENT:

Sulphur, as element ..... 90.0%

INERT INGREDIENT: ..... 10.0%

Sulphur - 90% through 325 mesh screen. TOTAL 100.0%

EPA Reg. No. 769-191 EPA Est. 769-AL-1

## CAUTION

### KEEP OUT OF REACH OF CHILDREN.

Avoid contamination of feed and foodstuffs. Do not use for medical purposes or as a feed supplement.

### ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**USE PRECAUTIONS:** Some fruits and other plants are susceptible to injury from sulphur under certain climatic conditions. The user is advised not to use sulphur on any crop unless local use has proven that sulphur is safe in that locality. During periods of high temperature, sulphur may burn foliage and fruit. Avoid applying this material during very hot periods of the day. When this product is used with Lead Arsenate, 8 pounds of hydrated lime or Basic Zinc Sulphate should be added for each 100 gallons of spray. Do not apply within 2 weeks of an oil spray treatment.

**CITRUS:** To control Rust Mite, apply 5 to 10 lbs. per 100 gallons of water. During cool weather use 10 lbs. of Security Wettable Sulphur per 100 gallons of water.

**APPLES:** To control Apple Scab and Powdery Mildew, apply 6 to 8 lbs. per 100 gallons of water. Make application between periods of rainfall. Repeat as necessary to cover new foliage growth during pink, petal-fall and cover spray periods.

**PEACHES and PLUMS:** To control Brown Rot and Scab, use 6 lbs. per 100 gallons of water. Follow locally issued spray schedule for number and frequency

of applications. Usually 6 applications are made. No. 1 - when  $\frac{2}{3}$  petals have fallen. No. 2 - when  $\frac{1}{4}$  shucks have shed. No. 3 - two weeks later. No. 4 - two weeks after No. 3. No. 5 - four weeks before harvest. No. 6 - two weeks before harvest. Do not apply within 2 weeks of an oil spray treatment.

### FLOWERS AND ORNAMENTALS

**APPLICATION:** To control the diseases and red spiders listed below, use Wettable Sulphur as a dust or spray using 6 pounds per 100 gallons of water. Applications should be made at 5 to 7 day intervals beginning when the disease first appears and continuing until all danger from the disease is past. During periods of humid or rainy weather, it may be necessary to make applications as often as every 2 to 3 days.

**POWDERY MILDEW:** Aster, Chrysanthemum, Cosmos, Dahlia, Daisy, Delphinium, Dogwood, Euonymus, Golden Fleece, Hibiscus, Honeysuckle, Hydrangea, Lady's Mantle, Ligustrum, Lilac, Crepe Myrtle, Phlox, Poplar, Silver Vine, Rose, Spiraea, Sunflower, Verbena, Willow and Zinna.

**LEAF SPOT:** Calendula, Cherry Laurel, Clematis, Columbine, English Ivy, Foxglove, Hydrangea, Petunia, Phlox, Sage, Smilax and Snapdragon.

**BLACK SPOT:** Rose.

**ROSE POWDERY MILDEW:** Rose.

**BOTRYTIS BLIGHT:** Hydrangea.

**RED SPIDERS:** Arborvitae, Carnation, Cedars, Chrysanthemum, Clematis, Hydrangea, Rose, Snapdragon and Sweetpeas.

### DISPOSAL WARNING

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

### NOTICE

**DISCLAIMER OF WARRANTY, MERCHANTABILITY AND LIABILITY** - If for any reason this product is not used, applied, stored, transported or disposed of in accordance with the instructions set forth on this label, then and thereafter Woolfolk Chemical Works, Inc. terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness.

Manufacturers

**WOOLFOLK CHEMICAL WORKS, INC.**

Atmore, Alabama

087130