NET WEIGHT 8 POUNDS

SECURITY

FERBAM

FUNGICIDE

ACTIVE INGREDIENT:
Ferbam (Ferric dimethyldithiocarbamate) 76.0%
INERT INGREDIENTS: 24.0%
Total Iron as metallic .... 10.2% TOTAL 100.0%
U.S.D.A. Reg. No. 769-215

ACCEPTED
3.6.1
8
769-215

CAUTION: KEEP OUT OF REACH OF CHILDREN. May cause irritation of eyes, nose, throat and skin! Avoid breathing dust or spray mist. Avoid contact with eyes, skin or clothing. In case of contact, flush with plenty of water; for eyes get medical attention. Keep away from fire and sparks.

PROTECT WILDLIFE: This product is toxic to fish. Keep out of any body of water.

DIRECTIONS

For specific directions on timing and rate of application consult local state or federal authorities or current spray schedules. Wash Ferbam into the spray tank through a deep screen or make a thick paste with a little water, then dilute with additional water and add directly to the spray tank while filling. THE FOLLOWING DILUTION RATES ARE BASED ON 100 GALLONS OF SPRAY.

FRUIT

APPLES: Apple Scab - Use 1½ to 2 pounds. Make applications at pre-bloom, calyx and the early cover sprays. Reduce to ¾ pound in late cover sprays. When used in combination with sulphur, use 1½ pound plus 3 pounds of ®NUTONEX sulphur. If russeting has been a problem, sulphur should not be used. Cedar Apple Rust - The above sprays plus one additional bloom spray at ½ pound will control this disease. Other Diseases - Use 1 to 1½ pounds for control of Frog-Eye Leaf Spot, Black Rot, Brooks Spot, Apple Blotch, Bitter Rot, Scab, Blotch, Fly Speck, and Blossom End Rot (Botrytis cinerea). Begin applications when diseases first appear and continue at 7 to 10-day intervals as long as diseases threaten. Do not use on Golden Delicious and other tender varieties as Ferbam may contribute to fruit russeting. Do not apply to apples within [illegible] harvest.

PEACHES: [illegible]

during the dormant period.

GRAPES: Black Rot - Use 2 pounds. Make application just before bloom, immediately after bloom and two weeks after bloom. Make additional applications at 10 to 14-day intervals as necessary. Do not apply within 7 days of harvest.

TOBACCO

Blue Mold: In plant beds - Use 3 to 4 pounds. Apply 3 gallons per 100 square yards when plants are small and increase to 6 gallons as plants are ready for transplanting. Make the first application when leaves are the size of a dime or when Blue Mold is first reported. Continue applications twice weekly until transplanting time.

PINE SEEDLINGS

Fusiform Rust in Plant Beds - Use 2 pounds at 3 to 5 day intervals from plant emergence to mid-June. Add suitable wetting agent to spray.

ORNAMENTALS

Use 1 to 1½ pounds to control Rust and Black Spot on ROSES; Septoria Leaf Spot of CHRYSANTHEMUMS; Rust on SNAPDRAGON; Bacterial Leaf Spot on GERANIUMS; Alternaria Blight and Rust of CARNATIONS; Botrytis Rot on CYCLAMEN and Botrytis Blight of TULIP. Begin applications when diseases first appear and continue at 7 to 10-day intervals as long as diseases threaten.

Camellia Flower Blight: Use 4 pounds as a drench at monthly intervals under directions of state quarantine officials.

DISPOSAL WARNING

[illegible]

NOTICE

[illegible]

WOOLFOLK CHEMICAL WORKS, LTD.

Manufacturers Fort Valley, Ga.