ACCEPTED
Mar 21, 1977
769-228
UNDER FOR
Commercial

SECURITY

7½% SEVIN® DUST

ACTIVE INGREDIENT:
Carbaryl (1-naphthyl N-methylcarbamate) 7.5%
INERT INGREDIENTS: 92.5%
® Trademark Union Carbide Corp. for 1-naphthyl N-methylcarbamate — TOTAL 100%

U.S.D.A. Reg. No. 769-228

CAUTION: KEEP OUT OF REACH OF CHILDREN.

May be absorbed through the skin. Do not breathe dust. Avoid contact with skin, eyes, or clothing. Avoid skin contact. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash clothing before reuse. Remove contaminated clothing and wash before reuse. Do not contaminate feed, foods, and water by storage or disposal.

PROTECT BEES

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your local Cooperative Agricultural Extension Service.

NOTE FOR PHYSICIAN

Carbaryl is a cholinesterase inhibitor. Atropine is antidotal.

DIRECTIONS

COTTON: To control Boll Weevil, Bollworm, Thrips, Flea Beetles, Cotton Fleahopper, Cotton Leafworm, Tarnished Plant Bug, Fall Armyworm, Cotton Leaf Perforator, Leaf Rollers, Leafhoppers, Grasshoppers, and for suppression of Aphids, apply at the rate of 15 to 25 pounds per acre. For control of Pink Bollworm, apply at the rate of 20 to 25 pounds per acre and for control of Salt-marsh Caterpillar and Stink Bugs, apply at 25 pounds per acre. Follow State recommendations for frequency and timing of applications. Usually, for early season control, 2 to 3 applications should be made at 7 to 14-day intervals at the rate of 15 to 20 pounds per acre. For mid season control, apply 20 to 25 pounds per acre. Repeat application at 5-day intervals as infestations are brought under control. Usually 4 applications will be necessary. Repeat a series of 4 applications at 5-day intervals if Weevils reappear in sufficient numbers to cause damage. For Thrips control, apply at the rate of 20 to 25 pounds per acre at 5-day intervals. Anytime Bollworms are a threat, when as many as 3 to 5 worms and eggs are found per 100 terminals examined, use 25 pounds of dust per acre. Direct the dust downward into the plants where the worms are hatching. Repeat applications at 5-day intervals until infestation is controlled. Allow seven days between last application and grazing treated fields or harvesting crop residue for use as feed or bedding for dairy or meat animals.

VEGETABLES

APPLICATION: Begin applications when insect or damage is first observed and repeat on a regular 5 to 7 day schedule or as necessary for good insect control.

BEANS: To control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetles, Leafhoppers, Flea Beetles, Japanese Beetles, and Corn Earworm - apply at the rate of 15 pounds per acre. To control Lygus Bug, Stink Bug and Lima Bean Pod Borer - apply 25 pounds per acre.

CABBAGE, COLLARDS, BROCCOLI, BRUSSEL SPROUTS, CAULIFLOWER, LETTUCE, MUSTARD GREENS, TURNIPS, KOHLRABI and SPINACH: To control Flea Beetle and Harlequin Bug - apply at the rate of 25 pounds per acre. To control Corn Earworm, Armyworm, Cabbage Caterpillars and small Cabbage Loopers - apply at 15 to 25 pounds per acre.

CUCUMBERS, SQUASH, CANTALOUPES, PUMPKINS, MELONS: To control Picklewarm, Melonworm, Cucumber Beetles, Squash Bug, Flea Beetles - apply at the rate of 12 pounds per acre.

CORN: To control Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetles, and Japanese Beetles - apply at the rate of 20 to 25 pounds per acre. For budworm damage in the whorl stage of corn, repeat as necessary. For maximum control of Corn Earworm, apply at 25 pounds per acre, preferably when silks are 1 to 2 days old. Applications after silks have turned brown will not be effective. Applications of Sevin dust to the tassels may seriously reduce the pollen shed.

CARROTS: To control Flea Beetles, Leafhoppers, Spittlebugs, Lygus and Stink Bugs - apply at the rate of 15 to 25 pounds per acre.

LETTUCE: To control Flea Beetles, Harlequin Bug, and Leafhoppers - apply at the rate of 15 pounds per acre. To control Corn Earworm, Armyworms, Cabbage Caterpillars, Spittlebugs, Lygus and Stink Bugs - apply at the rate of 15 to 25 pounds per acre.

OKRA: To control Corn Earworm and Stink Bugs - apply at the rate of 15 to 25 pounds per acre.

TOMATO, EGGPLANT, PEPPER: To control Tomato Fruitworm, Fall Armyworm, Flea Beetles and Leafhoppers - apply at the rate of 15 pounds per acre. For control of Tomato Hornworm, European Corn Borer and Stink Bugs - apply at the rate of 15 to 25 pounds per acre.

POTATO: To control Colorado Potato Beetle, Leafhoppers, and Flea Beetles, apply at 15 pounds and for Lygus Bug, apply at 25 pounds per acre.

INTERVAL BEFORE HARVEST: Allow 7 days between last application and harvest on Broccoli, Brussels Sprouts, Cabbage, Collards, Kohlrabi and Head Lettuce. Allow 14 days on Mustard Greens, Turnips and Spinach. On other vegetables listed on this label, application can be made up to and including the day of harvest.

FORAGE CROPS

APPLICATION: Apply when insects or signs of damage appear and repeat 7 to 14 days later if necessary.

ALFALFA, CLOVERS, PEANUTS, SOYBEANS, COWPEAS: To control Mexican Bean Beetle, Blister Beetles, Thrips, Leafhoppers, Three-Cornered Alfalfa Hopper, Cucumber Beetles, Bean Leaf Beetle, Japanese Beetle, Alfalfa Caterpillar and Velvet Bean Caterpillar, apply at the rate of 15 pounds per acre. To control Armyworms, Webworms, Corn Earworm, Stink Bugs, and Grasshoppers apply at the rate of 15 to 20 pounds per acre. Applications may be made up to and including the day of harvest. Some leaf curl may occur if applications are made when tender foliage is wet.

RICE, FORAGE GRASS, PASTURE: To control Thrips, Armyworms, Stink Bugs, and Grasshoppers - apply at the rate of 15 to 20 pounds per acre. For optimum thrips control in grasses grown for seed, direct dust into boot. Application may be made the day of harvest to grass and pasture. Allow 14 days between last application and harvest or graze.

CORN (Forage and Fodder): To control Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetles and Japanese Beetles, apply at the rate of 15 to 25 pounds per acre. Application may be made up to within 7 days of harvest.

SORGHUM (Grain and Forage): To control Armyworms, Webworms, Corn Earworm, Stink Bugs, and Grasshoppers, apply at the rate of 15 to 25 pounds per acre. Allow 21 days between last application and harvest of grain, no time limit on sorghum forage.

TOBACCO

To control Hornworms, Budworms and Flea Beetles, apply at the rate of 25 to 35 pounds per acre. Begin applications when worms are small and repeat as necessary for control. Avoid excessive applications. For budworm control with a hand duster, direct the dust to the bud and upper leaves of each plant, only 5 to 6 pounds per acre is necessary. Do not apply with cloth bags or apply heavily because plant injury may result.

DISPOSAL WARNING

Dispose of waste dust in a sanitary landfill or by burying it in areas away from water supplies. Do not contaminate water by disposing of wastes or by burning.

NOTICE

The seller makes no warranties [illegible] 176728

ACCEPTED WITH COMMENTS