

SECURITY

10% SEVIN[®] DUST

Jul 21 1954
THE FEDERAL INSECTICIDE
AND RODENTICIDE ACT

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) 10.0%

INERT INGREDIENTS: 90.0%

U. S. D. A. Reg. No. 769-229 TOTAL 100.0%

(R) Trademark Union Carbide Corp. for 1-naphthyl N-methylcarbamate

CAUTION! KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. SKIN CONTACT MAY BE HARMFUL. Avoid contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

PROTECT BEES: This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

AVOID CONTAMINATION OF FOODSTUFFS: Avoid contamination of feed, feeding troughs, and watering receptacles.

DIRECTIONS

COTTON: To control Boll Weevil, Bollworm, Thrips, Flea Beetles, Cotton Fleahopper, Cotton Leafworm, Tarnished Plant Bug, Fall Armyworm, Cotton Leaf Perforator, Leaf Rollers, Leafhoppers, Grasshoppers, and for suppression of Aphids, apply at the rate of 10 to 20 pounds per acre. For control of Pink Bollworm, apply at the rate of 15 to 25 pounds per acre, and for control of Salt-marsh Caterpillar and Stink Bugs, apply at 20 pounds per acre. Follow State recommendations for frequency and timing of applications. Usually, for early season control, 2 to 3 applications should be made at 7 to 10-day intervals at the rate of 8 to 10 pounds per acre. For mid season control, apply 10 to 15 pounds per acre. Repeat application at 5-day intervals until infestation is brought under control. Usually 4 applications will be necessary. Repeat a series of 4 applications at 5-day intervals if boll weevils reappear in sufficient numbers to cause damage. For late season control, apply at the rate of 15 to 20 pounds per acre at 5-day intervals. Anytime bollworms are a threat, when as many as 4 to 5 small worms and eggs are found per 100 terminals examined, apply at least 15 pounds of dust per acre. Direct the dust downward into the tops of the plants where the worms are hatching. Repeat applications at 5-day intervals until infestation is controlled. Allow seven days between last application and grazing treated fields or harvesting crop residue for use as feed or bedding for dairy or meat animals.

VEGETABLES

APPLICATION: Begin applications when insect or damage is first

observed and repeat on a regular 5 to 7 day schedule or as necessary for good insect control.

BEANS: To control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetles, Leafhoppers, Flea Beetles, Japanese Beetles, and Corn Earworm - apply at the rate of 10 pounds per acre. To control Lygus Bug, Stink Bug and Lima Bean Pod Borer - apply 20 pounds per acre.

CABBAGE, COLLARDS, BROCCOLI, BRUSSEL SPROUTS, CAULIFLOWER, LETTUCE, MUSTARD GREENS, TURNIPS, KOHLRABI and SPINACH: To control Flea Beetle and Harlequin Bug - apply at the rate of 10 pounds per acre. To control Corn Earworm, Armyworm, Cabbage Worm and small Cabbage Loopers - apply 10 to 20 pounds per acre.

CUCUMBERS, SQUASH, CANTALOUPE, PUMPKINS, MELONS: To control Pickleworm, Melonworm, Cucumber Beetles, Squash Bug, Flea Beetles - apply at the rate of 10 pounds per acre.

CORN: To control Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetles, and Japanese Beetles - apply at the rate of 15 to 20 pounds per acre. For budworm damage in the whorl, apply to entire plant and repeat as necessary. For insects attacking silks and ears, apply at 2 to 3-day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending upon severity of infestation.

CARROTS: To control Flea Beetles, Leafhoppers, Spittlebugs, Lygus and Stink Bugs - apply at the rate of 10 to 20 pounds per acre.

LETTUCE: To control Flea Beetles, Harlequin Bug, and Leafhoppers - apply at the rate of 10 pounds per acre. To control Corn Earworm, Armyworms, Cabbage Caterpillars, Spittlebugs, Lygus and Stink Bugs - apply at the rate of 10 to 20 pounds per acre.

OKRA: To control Corn Earworm and Stink Bugs - apply at the rate of 10 to 20 pounds per acre.

TOMATO, EGGPLANT, PEPPER: To control Tomato Fruitworm, Fall Armyworm, Flea Beetles and Leafhoppers - apply at the rate of 10 pounds per acre. For control of Tomato Hornworm, European Corn Borer and Stink Bugs - apply at the rate of 10 to 20 pounds per acre.

POTATO: To control Colorado Potato Beetle, Leafhoppers, and Flea Beetles, apply at 10 pounds and for Lygus Bug, apply at 20 pounds per acre.

INTERVAL BEFORE HARVEST: Allow 3 days between application and harvest on Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kohlrabi and Head Lettuce. Allow 14 days on Collards, Leaf Lettuce, Mustard Greens and Turnips. On other vegetables listed on this label application can be made up to and including the day of harvest.

FORAGE CROPS

APPLICATION: Apply when insects or signs of damage appear and repeat 7 to 14 days later if necessary.

ALFALFA, CLOVERS, PEANUTS, SOYBEANS, COW-PEAS: To control Mexican Bean Beetle, Blister Beetles, Thrips, Leafhoppers, Three-Cornered Alfalfa Hopper, Cucumber Beetles, Bean Leaf Beetle, Japanese Beetle, Alfalfa Caterpillar, and Velvet Bean Caterpillar, apply at the rate of 10 pounds per acre. To control Armyworms, Webworms, Corn Earworm, Stink Bugs, and Grasshoppers, apply at the rate of 10 to 15 pounds per acre. Application may be made up to and including the day of harvest. Some leaf injury may occur if applications are made when tender foliage is wet.

RICE, FORAGE GRASS, PASTURE: To control Thrips, Armyworms, Stink Bugs, and Grasshoppers - apply at the rate of 10 to 15 pounds per acre. For optimum thrips control in grasses grown for seed, direct dust into boot. Application may be made the day of harvest to grass and pasture. Allow 14 days between last application and harvest on rice.

CORN (Forage and Fodder): To control Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetles and Japanese Beetles, apply at the rate of 10 to 20 pounds per acre. Application may be made up to within 7 days of harvest.

SORGHUM (Grain and Forage): To control Armyworms, Webworms, Corn Earworm, Stink Bugs, and Grasshoppers, apply at the rate of 10 to 20 pounds per acre. Allow 21 days between last application and harvest of grain, no time limit on sorghum forage.

TOBACCO

To control Hornworms, Budworms, and Flea Beetles apply at the rate of 15 to 25 pounds per acre. Begin applications when worms are small and repeat as necessary for control. Avoid excessive applications. For budworm control with a hand duster, direct the dust to the bud and upper leaves of each plant, only 4 to 6 pounds per acre is necessary. Do not apply with cloth bags or apply heavily because plant injury may result.

DISPOSAL WARNING

Dispose of waste pesticide and container by burying in non-crop land away from water supplies. Never re-use container.

NOTICE

DISCLAIMER OF WARRANTY, MERCHANTABILITY AND LIABILITY

If for any reason this product is not used, applied, stored, transported or disposed of in accordance with the instructions set forth on this label, then and thereafter Woolfolk Chemical Works, Ltd. terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness. 17042

Manufacturers

WOOLFOLK CHEMICAL WORKS, LTD.

East Valley, Ga.