

ACCEPTED  
 JUN 28 1985  
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 769-254

reg # 769-254  
 11217  
 1041

necessary near and during harvest if conditions are favorable for brown rot development. If a mite problem develops following the use of this product, apply a good miticide. Consult us or your state agricultural experiment station for additional information.

**POULTRY**

CHICKENS, TURKEYS, DUCKS, GEESE, GAME BIRDS AND PIGEONS  
 Control chicken mite, lice, fleas and bedbugs. Use conventional power spray equipment when spraying buildings for control of these pests. Use 8 pounds of Sevin-Sulphur 2-6 Spray per 25 gallons of water. Spray 1 to 2 gallons per 1000 square feet of wall bedding, litter or roost surface. Force spray into cracks and crevices. Repeat as needed. Ventilate while spraying. Do not use within 7 days of slaughter. Avoid contamination of nests, eggs and feeding and watering troughs.

**STORAGE AND DISPOSAL**

**STORAGE:** Store pesticide in original container and place in a locked storage area. Opened pesticide containers should be securely resealed prior to storage. Protect from penetration of moisture, excessive heat or cold. Avoid cross-contamination with other pesticides or fertilizers. Use caution when storing pesticides in stacks.

**PESTICIDE DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment, then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**NOTICE**

**DISCLAIMER OF WARRANTY, MERCHANTABILITY AND LIABILITY:** If for any reason this product is not used, applied, stored, transported or disposed of in accordance with the instructions set forth on this label, then and thereafter Security Chemical Company terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness.

NET WEIGHT: 10 POUNDS

**SECURITY CHEMICAL COMPANY**  
 Manufacturers Atlanta, GA



**Sevin-Sulphur  
 2-6 Spray**

**ACTIVE INGREDIENTS:**

Carbaryl (1-naphthyl N-methylcarbamate)	12.50%
Sulphur, elemental	69.75%
<b>INERT INGREDIENTS:</b>	<b>17.75%</b>
	<b>TOTAL 100.00%</b>

EPA REG NO 769-254      EPA EST 769-GA-1  
 \* Trademark Union Carbide Corporation for 1-naphthyl N-methylcarbamate

**KEEP OUT OF REACH OF CHILDREN  
 CAUTION**

(See additional precautionary statements on the side panel.)

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.  
**IF INHALED:** Remove victim to fresh air. **IF ON SKIN:** Wash with soap and water. Remove any contaminated clothing. **IF IN EYES:** Flush eyes with water.  
**NOTE TO PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates or cholinesterase-inhibiting drugs.

**PRECAUTIONARY STATEMENTS**

**CAUTION.** May be harmful if swallowed. Avoid breathing vapors (dust or spray mist). Avoid contact with eyes, skin or

clothing. Wash thoroughly after handling.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of waste. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

**PEACHES.** To control curculio, Oriental fruit moth, cat-facing insects, brown rot and scab, apply this product at the rate of 8 pounds per 100 gallons of water. Applications should not be made when temperatures is excessively high because of the danger of sulphur injury. The following schedule is suggested as a general guide but is in no way guaranteed. The number of applications and times of application may vary in different localities. Early ripening varieties will not require the full schedule of applications. Consult your state agricultural experiment station or state spray schedule for specific directions.

DO NOT APPLY THIS MATERIAL WITHIN 1 DAY OF HARVEST

**SPRAY SCHEDULE**

- 1 When all petals are off
- 2 When 1/4 shucks are shed
- 3 7 to 10 days after No. 2
- 4 12 to 14 days after No. 3
- 5 14 days after No. 4 (see note below)
- 6 6 weeks before harvest of each variety
- 7 4 weeks before harvest
- 8 2 weeks before harvest
- 9 1 week before harvest (see note below)

**NOTE:** Unless curculios or other insects are present at the time of sprays 5 and 9, this material can be omitted and only a Nutonex Sulphur spray for control of brown rot and scab need be applied. Additional sulphur or other fungicidal sprays or dusts may be