

[769-256]

to deliver a fine mist, and apply this material at the rate of 1 pint to 8,000 cubic feet. Direct spray toward and into all places commonly inhabited by the insects. Move sprayer about the room to thoroughly fill the area with fog or vapor. As a **SURFACE SPRAY** - to control roaches, silverfish, crickets, spiders, and ants in the above areas - use a good sprayer, adjusted to deliver a coarse droplet spray, at the rate of 1 gallon to 750 square feet of surface. Application should be made to cracks, crevices, recesses, and other hiding and breeding areas. Whenever possible, keep doors and windows closed for 10 minutes or longer after application. Sweep up insects and burn. When using in food processing plants, food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when facility is not in operation.

TO CONTROL exposed stages of clothes moths, carpet beetles, fleas, brown dog ticks, roaches, silverfish, crickets, spiders, and ants in hotels, motels, homes, and other buildings, use as directed above for surface spray.

TO CONTROL exposed stages of cheese skipper and cheese mites in restaurants, homes and other areas where they may occur - use as directed above for surface spray.

TO CONTROL clothes moths - use a good sprayer, and spray all infested fabrics thoroughly, particularly the folds and seams. Unless treated fabrics are in moth-tight containers, repeat applications every 30 days.

TO CONTROL bedbugs - use a good sprayer and spray sleeping quarters of humans and animals. Thoroughly spray mattresses, beds, and other surfaces where bedbugs congregate or hide. Apply at the rate of 1 gallon to 750 square feet of surface.

TO CONTROL fleas and brown dog ticks - thoroughly spray infested rugs and carpets, clothes storage area and clothing. Spray liberally in and around animals' sleeping quarters, kennels and dog houses, forcing spray into cracks and crevices. Repeat application every 30 days unless treated clothing is stored in tight containers.

NOTICE: The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

NET CONTENTS

SECURITY

ONE QUART

MILL SPRAY

ACTIVE INGREDIENTS:	100.00%
Technical Piperonyl Butoxide*	1.50%
Pyrethrins15%
Petroleum Hydrocarbons	98.35%

*Equivalent to 1.2% of (Butylcarbityl) (6-propylpiperonyl) ether and .30% of related compounds.

E. P. A. Reg. No. 769-256

CAUTION! KEEP OUT OF REACH OF CHILDREN. See side panel for additional precautions.

PROTECT WILDLIFE: This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of waste. Apply this product only as specified on the label.

DISPOSAL WARNING: Do not reuse container. Destroy when empty.

CAN BE USED IN: Mills, elevators, granaries, warehouses, food storages, seed stores, food processing plants, restaurants, hotels, motels, homes, and other buildings. 271019

WOOLFOLK CHEMICAL WORKS, LTD.
Manufacturers Fort Valley, Ga.

CAUTION: Harmful if Swallowed! Do not use or store near heat or open flame. To avoid any explosion hazard, when using vaporizers, not more than 1 gallon of material should be atomized in each 40,000 cubic feet. Pilot lights should be extinguished, an important precautionary measure when sprays are used in poorly ventilated rooms. Remove pets, birds, and cover fish aquariums before spraying. When using in enclosed areas, do not remain in treated area. Ventilate the area after treatment.

DIRECTIONS

Before spraying in mills, elevators, and granaries, clean up waste grain, dust, dirt, and debris. Treat unloading, handling and processing areas, and apply to conveyors, hoppers, and other machinery and equipment.

TO CONTROL flies, mosquitoes, gnats, wasps and sand flies in mills, elevators, granaries, warehouses, food storages, food processing plants, restaurants, hotels, motels, homes, and other buildings, apply as follows: As a **SPACE** or **VAPOR SPRAY** use a good electric, steam or compressed air applicator, adjusted to deliver a fine mist, and apply this material at the rate of 1 pint to 8,000 cubic feet. Direct the spray toward the upper portions of the area being treated and move the sprayer about the room to thoroughly fill the area with fog or vapor. As a **SURFACE SPRAY** apply with a good sprayer, adjusted to deliver a coarse droplet spray, at the rate of 1 gallon to 750 square feet of surface. Application should be made to the following resting places of pests: screens, doors, around lights and ceilings. Whenever possible, keep doors and windows closed for 10 minutes or longer after application. Sweep up insects and burn. When using in food processing plants, foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when facility is not in operation.

TO CONTROL exposed stages of the following grain and stored products infesting pests: saw toothed grain beetle, angoumois grain moth, Mediterranean flour moth, meal worms, cigarette beetle, grain mites, in mills, elevators, granaries, warehouses, food storages, food processing plants, restaurants, homes, and other buildings, apply as follows: As a **SPACE** or **VAPOR SPRAY** - use a good electric, steam or compressed air applicator, adjusted

ACCEPTED
NOV 27, 1972
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 769-256 SUBJECT
TO APPLICABLE ORDINANCES.