



000769-00271-100462

Printed: 11:12:35 Tuesday, 18 Jul, 2000 # 10 / 9055

Systems Integration Group, Inc.

769-271

10/04/1962

1/2

SECURITY BRAND CHEMICALS • SECURITY BRAND CHEMICALS •
 MICALS • SECURITY BRAND CHEMICALS • SECURITY BRAND
 URITY BRAND CHEMICALS • SECURITY BRAND CHEMICALS •
 MICALS • SECURITY BRAND CHEMICALS • SECURITY BRAND
 URITY BRAND CHEMICALS • SECURITY BRAND CHEMICALS •
 MICALS • SECURITY BRAND CHEMICALS • SECURITY BRAND
 URITY BRAND CHEMICALS • SECURITY BRAND CHEMICALS •
 MICALS • SECURITY BRAND CHEMICALS • SECURITY BRAND

ORNAMENTALS

TURF, LAWNS: To control—Chinch bugs, Fall armyworm, Leafhoppers, Sod webworm and Lawn moths—apply at the rate of 1 cupful in 15 to 20 gallons of water to 500 square feet. Apply after rain or watering and do not water for at least 2 days after application. Repeat 2 to 3 weeks later if necessary.

FLOWERS, SHRUBS, TREES: To control—Apple aphid, Bagworm, Birch leaf miner, Blister beetles, Boxelder bug, Boxwood leaf miner, Elm leaf aphid, Elm leaf beetle, Flea beetles, Japanese beetle, Lace bugs, Leafhoppers, Leaf rollers, Oak leaf miner, Orange tortrix, Periodical cicada, Plant bugs, Psyllids, Rose aphid, Rose-slug, Scale insects, Tent caterpillars, Willow leaf beetle and exposed Thrips—use 2 pounds per 100 gallons or 2½ level tablespoonfuls per gallon. Apply when insect or damage first appears and repeat at weekly intervals or as needed. For scale insect control apply in Spring and early Summer when scale crawlers are present.

TOBACCO

To control Hornworms, Budworms and Flea Beetles, apply at the rate of 2 to 4 pounds per acre. Treated fields may be entered immediately after foliage has dried. If late applications are necessary treat immediately after priming rather than before priming.

POULTRY

POULTRY HOUSES: For use in and around poultry houses, to control Chicken mite, Fleas and Bed bugs—use 2 pounds per 25 gallons of water. For Fowl Ticks use 8 pounds per 25 gallons of water. Spray 1 to 2 gallons per 1000 square feet of wall, bedding, litter or roost surfaces. Force spray into cracks and crevices. Ventilate while spraying. Repeat as needed. Do not treat nest litter. Avoid direct contamination of eggs. Do not use on poultry or game birds. For application to birds use Security 5% Sevin Dust. Do not use in premise treatments within 7 days of slaughter.

SECURITY

BRAND

NET WEIGHT TWO POUNDS

50%

SEVIN*

WETTABLE

FOR USE ON

FRUIT, VEGETABLES, ORNAMENTALS

ACTIVE INGREDIENT

1-naphthyl N-Methylcarbamate 50.00%

INERT INGREDIENTS 50.00%

Total 100.00%

CAUTION: Harmful if Swallowed or Inhaled. Do not take internally. Avoid breathing dust or spray mist. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs.

*Trademark of Union Carbide Corp. for 1-Naphthyl N-Methylcarbamate

WOOLFOLK

CHEMICAL WORKS, LTD.

MANUFACTURERS • FORT VALLEY, GA

CURITY BRAND CHEMICALS • SEC

CHEMICALS • SECURITY BRAND

CURITY BRAND CHEMICALS •

CHEMICALS • SECURITY BRAN

CURITY BRAND CHEMICALS •

EMIC

Y BRAN

NOTE

COMPATIBILITY: Sevin is compatible with commonly used insecticides such as lead arsenate, DDT and other chlorinated hydrocarbons, parathion, and other organic phosphates, and botanicals; also with such fungicides as glyodin, organomercury compounds, fixed coppers, dithiocarbamates, sulfur, captan, Cyprex, and Phallon. Sevin is unstable under highly alkaline conditions. Not effective if used with alkaline materials such as Bordeaux, lime, Eme sulfur and caseinlime spreaders.

PHYTOTOXICITY: Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following spraying. May cause foliar injury when used before second cover on McIntosh and York apples in the eastern United States. Foliar injury may result from combination with summer oils on apples and pears in the Pacific Northwest.

Sevin does not control mites but is compatible with all common miticides. It is toxic to honeybees. Avoid use on crop plants during flowering and pollination period.

NOTICE

Woolfolk Chemical Works, Ltd., makes no warranty, express or implied concerning this material except that it conforms to the chemical description on the label. Neither Woolfolk Chemical Works, Ltd., nor the seller shall be held responsible in any manner for any personal injury or property damage or other types of loss resulting from the handling, storage or use of this material. The buyer assumes all risk and liability therefrom and accepts and uses this material on these conditions.

SECURITY
BRAND

DIRECTIONS

FRUIT

Dosages recommended refer to pounds of 50% SEVIN per 100 gallons of water. Apply for full coverage in normal spray schedules. For scale control apply when crawlers are present in Spring and early Summer. To control rosy aphid apply before leaves are curled. Apply at 10-day intervals for optimum apple maggot control. For lesser peach tree borer control spray limbs and trunk thoroughly.

APPLES, PEARS: To control—Codling moth, Green apple aphid, White apple leafhopper and Apple mealy-bug—use 1 pound. Use 2 pounds to control Apple Maggot, Plum Curculio, Rosy apple aphid, Red-banded leaf roller, Fruit tree leaf roller, Pear psylla, Periodical cicada, Eyespotted bud moth, European apple sawfly, Eastern tent caterpillar, Japanese beetle, Tentiform leaf miner, Forbes scale, San Jose scale, Oystershell scale, and Lecanium scale. Application of Sevin within 30 days after full bloom may cause fruit thinning. To avoid this, delay use until at least 30 days after bloom.

PEACHES: To control—Plum curculio, Oriental fruit moth, Peach twig borer, lesser Peach tree borer, Cat-facing plant bugs, Periodical cicada, Japanese beetle, San Jose scale, Lecanium scale, Olive scale and Forbes scale—use 2 pounds.

PLUMS, PRUNES, CHERRIES: To control—Cherry fruit fly, Fruit tree leaf roller, Peach twig borer, lesser peach tree borer, Black cherry aphid, Eyespotted bud moth, Plum curculio, Red-banded leaf roller, Japanese beetle, Forbes scale, San Jose scale, Oystershell scale, and Lecanium scale—use 2 pounds. Use 1½ pounds to control Codling moth, Eastern Tent caterpillar, Orange tortrix and Western tussock moth.

GRAPES: To control—Grape leaf folder and Grape leafhopper—use 1 to 2 pounds. Use 2 pounds for Grape berry moth and Japanese beetle. Apply at the rate of 200 gallons dilute spray per acre. Apply just before first broad larvae emerge from rolls. Apply as needed for leafhoppers.

STRAWBERRIES: To control—Meadow spittlebug and Strawberry leaf roller—use 2 pounds and apply at 200 gallons per acre.

INTERVAL: This material may be applied up to within 1 day of harvest on Apples, Pears, Peaches, Plums, Prunes, Cherries and Strawberries. It may be applied the day of harvest on Grapes.

VEGETABLES

Rates recommended refer to pounds of 50% SEVIN per acre. Use the amount of water necessary to provide adequate coverage. Begin applications when insect or damage appears and continue at 7 to 10 day intervals or as necessary for good control.

BEANS: To control—Mexican bean beetle—use 1 pound. Use 2 pounds to control Bean leaf beetle, Cucumber beetle, Western bean cutworm, Flea beetle and Japanese beetle. Use 2 to 3 pounds to control Corn earworm. Armyworms and Velvetbean caterpillar. Use 4 pounds to control Lima bean pod borer, Lygus bug and Stink bug.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI: To control—Flea Beetle and Harlequin bug apply at the rate of 1 to 2 pounds per acre. To control—Corn earworm, Armyworm and Cabbage caterpillars and small Cabbage loopers, apply at the rate of 2 to 4 pounds per acre.

CORN: To control—Corn earworm, European corn borer, Fall armyworm, Sap beetles and Japanese beetle—use 2 to 4 pounds. Where budworm damage is occurring apply to entire plant, directing the spray downward into the bud. Repeat as necessary. For corn earworm begin when first silks appear and direct to the zone of ear production. Repeat applications at 2 to 3 day intervals until silks begin to dry. Three or more applications may be required depending upon the severity of infestation. Allow 7 days between last application and harvest if corn is to be used as forage or fodder to be fed dairy or meat animals.

CUCUMBERS, SQUASH, CANTALOUPE, PUMPKINS, MELONS: To control—Pickworm, Melonworm, Cucumber beetles, Squash bug, Flea beetles, apply at the rate of 2 pounds per acre.

CARROTS: To control—Flea beetles, Leafhoppers, Spittlebugs and Stink bugs, apply at the rate of 1 to 2 pounds per acre.

LETTUCE: To control—Flea beetles, Harlequin bug, and Leafhoppers, apply at the rate of 1 to 2 pounds per acre. To control—Corn earworm, Armyworms, Cabbage caterpillars, Spittlebugs, Lygus and Stink bug, apply at the rate of 2 to 4 pounds per acre.

OKRA: To control—Corn earworm and Stink bugs, apply at the rate of 2 to 4 pounds per acre.

POTATO: To control—Colorado potato beetle—use 1 pound. Use 2 pounds to control Leafhoppers and Flea Beetles. Use 4 pounds to control Lygus bug.

TOMATO, EGGPLANT, PEPPER: To control—Tomato fruitworm, Fall armyworm, Flea beetles and Leafhoppers—use 2 pounds. Use 2 to 4 pounds to control Tomato hornworm, European corn borer and Stink bugs.

INTERVAL BEFORE HARVEST: This product can be used up to and including the day of harvest on all vegetables listed except Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kohlrabi and Lettuce. When applied to these vegetables allow 3 days between the last application and harvest.

ADDITIONAL DIRECTIONS: "See side panel for use on ornamentals, tobacco and poultry."

WOOLFOLK
CHEMICAL WORKS, LTD.
MANUFACTURERS • FORT VALLEY, GA.