



000769-00271-021263

Printed: 11:13:03 Tuesday, 18 Jul, 2000 # 12 / 9057

Systems Integration Group, Inc.

WOOLFOLK CHEMICAL WORKS, LTD.



FARM AND ORCHARD CHEMICALS

FORT VALLEY, GEORGIA

769-271

ACCEPTED
 Feb 12-1963
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO.



LTD., FORT VALLEY, GEORGIA

50% SEVIN*

WETTABLE

LTD., FORT VALLEY, GEORGIA

50% SEVIN* WETTABLE

DIRECTIONS: Read label carefully and use only in accordance with directions. To avoid injury to young tender foliage avoid application during periods of high humidity or when plants are wet and when plants are suffering from drought. Use 2½ level tablespoonfuls per gallon of water (4 oz. to 12½ gallons of water) and apply thoroughly. For scale insect control apply when crawlers are present in Spring and early Summer. To avoid killing honeybees avoid applications to plants during flowering and pollination period.

ORNAMENTALS, FLOWERS, SHRUBS, TREES: To control Aphids, Bagworm, Flea beetles, Japanese beetle, Lace bugs, Leafhoppers, Leaf miners, Leaf rollers, Periodical cicada (17-year locust), Plant bugs, Rose-slug, Scale insects, Tent caterpillars and exposed Thrips, apply when insect or damage first appears and repeat at weekly intervals or as needed. Do not apply to Boston Ivy.

FRUIT—APPLES, PEARS, PEACHES, PLUMS, CHERRIES, GRAPES: To control Codling moth, Apple maggot, Plum curculio, Leaf rollers, Oriental fruit moth, San Jose scale, Japanese beetle, apply at petal-fall and at 10 to 14-day intervals as necessary during the season. To

avoid fruit thinning do not apply to apples until 30 days after full bloom. May cause injury to McIntosh and York apples if applied within 15 days of petal fall. To control Rosy apple aphid apply before leaves are curled. Applications can be made up to within 1 day of harvest.

VEGETABLES—BEANS, CABBAGE, CORN, CUCUMBERS, SQUASH, LETTUCE, POTATO, TOMATO, EGGPLANT, PEPPER: To control Mexican bean beetle, Flea beetles, Harlequin bug, Cucumber beetle, Corn earworm, Armyworm, Cabbage caterpillars, small Cabbage loopers, Pickle-worm, Mellow-worm, Stink bugs, Colorado potato beetle, Tomato fruitworm, apply when insects or damage appears and repeat at 7 to 10-day intervals as necessary. Can be applied up to within 3 days of harvest of Cabbage and Lettuce and to within 1 day on all other vegetables listed except corn to be harvested for storage where the interval should be 7 days.

LAWN INSECTS: To control Chinch bugs, Fall armyworm, Leafhoppers, Sod webworm and Lawn moths, apply at the rate of 4 oz. in 15 to 20 gallons of water to 500 square feet. Apply after rain or watering and do not water for at least 2 days after application. Repeat 2 to 3 weeks later if necessary.

*Trademark of Union Carbide Corp. for 1-Naphthyl, N-Methylcarbamate

FOR FRUITS
 FLOWERS
 VEGETABLES
 ORNAMENTALS
 & LAWNS



ACTIVE INGREDIENT

Carbaryl (1-naphthyl N-Methylcarbamate)	50.0%
INERT INGREDIENTS	50.0%
TOTAL	100%

CAUTION: Harmful if swallowed. Avoid prolonged breathing of dust or spray mist. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid storage near feed and food products. Keep out of reach of children.

NOTICE: Woolfolk Chemical Works, Ltd., makes no warranty, express or implied concerning this material except that it conforms to the chemical description on the label. Neither Woolfolk Chemical Works, Ltd., nor the seller shall be held responsible in any manner for any personal injury or property damage or other types of loss resulting from the handling, storage or use of this material. Buyer assumes all risk and liability therefrom and accepts and uses this material on these conditions.