



000769-00271-041067

Printed: 11:13:26 Tuesday, 18 Jul, 2000 # 14 / 9059

Systems Integration Group, Inc.

SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY

LTD., FORT VALLEY, GEORGIA

50% SEVIN*

WETTABLE

LTD., FORT VALLEY, GEORGIA



50% SEVIN* WETTABLE

DIRECTIONS: Read label carefully and use only in accordance with directions. To avoid injury to young tender foliage avoid application during periods of high humidity or when plants are wet and when plants are suffering from drought. Use 2½ level tablespoonfuls per gallon of water (4 oz. to 12½ gallons of water) and apply thoroughly. For scale insect control apply when crawlers are present in Spring and early Summer. To avoid killing honeybees avoid applications to plants during flowering and pollination period.

ORNAMENTALS, FLOWERS, SHRUBS, TREES: To control Aphids, Bagworm, Flea beetles, Japanese beetle, Lace bugs, Leafhoppers, Leaf miners, Leaf rollers, Periodical cicada (17-year locust), Plant bugs, Rose-slug, Scale insects, Tent caterpillars and exposed Thrips, apply when insect or damage first appears and repeat at weekly intervals or as needed. Do not apply to Boston Ivy.

FRUIT—APPLES, PEARS, PEACHES, PLUMS, CHERRIES, GRAPES: To control Codling moth, Apple maggot, Plum curculio, Leaf rollers, Oriental fruit moth, San Jose scale, Japanese beetle, apply at petal-fall and at 10 to 14-day intervals as necessary during the season. To

avoid fruit thinning do not apply to apples until 30 days after full bloom. May cause injury to McIntosh and York apples if applied within 15 days of petal fall. To control Rosy apple aphid apply before leaves are curled. Applications can be made up to within 1 day of harvest.

VEGETABLES—BEANS, CABBAGE, CORN, CUCUMBERS, SQUASH, LETTUCE, POTATO, TOMATO, EGGPLANT, PEPPER: To control Mexican beetle, Flea beetles, Harlequin bug, Cucumber caterpillars, small Cabbage loopers, Pickworm, Mellow-worm, Stink bugs, Colorado potato beetle, Tomato fruitworm, apply when insects or damage appears and repeat at 7 to 10-day intervals as necessary. Can be applied up to within 3 days of harvest of Cabbage and Lettuce and to within 1 day on all other vegetables listed except corn to be harvested forage where the interval should be 7 days.

LAWN INSECTS: To control Chinch bugs, Flea beetle, Armyworm, Leafhoppers, Sod webworm and Lawn moths, apply at the rate of 4 oz. in 10 to 20 gallons of water to 500 square feet. Apply after rain or watering and do not water for at least 2 days after application. Repeat 2 to 3 weeks later if necessary.

*Trademark of Union Carbide Corp. for 1-Naphthyl, N-Methylcarbamate

ACTIVE INGREDIENT

Carbaryl (1-naphthyl N-Methylcarbamate)	50.0%
INERT INGREDIENTS	50.0%
USDA Reg. No. 769-271	TOTAL 100%

CAUTION: Harmful if inhaled, swallowed, or absorbed through the skin. Do not breathe the dust. Do not take internally. Avoid skin contact. Avoid getting in eyes. Wash hands before eating. Wash

all contaminated clothing with soap and water before re-use. Avoid contamination of foodstuffs, feed, troughs, and water receptacles.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

NOTICE: Woolfolk Chemical Works, Ltd., makes no warranty, express or implied concerning this material except that it conforms to the chemical description on the label.

ACCEPTED
Open-10-1967
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 769-271

769-271

04/10/1967

1/1