days of harvest. Do not feed treated threshings to livestock or allow livestock to graze in treated fields. Do not use on lima beans.

BROCCOLI, CABBAGE, CAULIFLOWER: To control - Cabbage Looper, Imported Cabbage Worm, Diamond-back Moth Larvae, Cross-striped Cabbage Worm, Flea Beetle, Harlequin Bug and Stink Bug - apply at the rate of 1½ quarts per acre. Do not apply after edible parts begin to form.

EGGPLANT, PEPPERS: To control - Colorado Potato Beetle, Flea Beetle, Whitefly and Aphids - apply at the rate of 1 quart per acre. Apply when insects are first observed and repeat as necessary for control. Do not apply to eggplant or peppers within 1 day of harvest.

POTATOES: To control - Flea Beetle, Colorado Potato Beetle, Leafhopper and Aphids - apply at the rate of 1 quart per acre. To control - Southern Armyworm, Green Stink Bug, Plant Bugs, Potato Tuberworm and Whitefly - apply at the rate of 1 to 2 quarts per acre. Begin applications when insects are first observed and repeat as necessary for control. Do not plant root crops other than carrots, potatoes and sweet potatoes as follow-up crops.

TOMATOES: To control - Aphids, Colorado Potato Beetle, Flea Beetle, Green Stink Bug and Blister Beetle - apply at the rate of 1 quart per acre. Do not apply within 1 day of harvest. When applying in greenhouse wear mask or respirator of a type passed by the U.S. Bureau of Mines for Endosulfan

SMALL FRUITS

STRAWBERRIES: [illegible] White, Western Cabbage Looper, Strawberry Aphid and [illegible] quart per [illegible] water or Dasheen at the rate of 4 gallons per acre for [illegible] Apply [illegible] gallons of spray [illegible] Caution: [illegible]

The seller makes no express warranty of merchantability [illegible] other than those which are stated beyond the description on this label.

ACCEPTED

CONTENTS

SECURITY

ONE GALLON

ENDOSULFAN

EMULSIFIABLE CONCENTRATE

TWO POUNDS PER GALLON

ACTIVE INGREDIENTS:

Endosulfan Hexachlorohexahydr[illegible] [illegible]

Heavy Aromatic Solvent [illegible]

INERT INGREDIENTS: [illegible]

E. P. A. Reg. No. [illegible]

Reg. Endosulfan Sales Stocks [illegible]

KEEP OUT OF REACH OF CHILDREN

[illegible] to be immediately removed [illegible] Water [illegible] Water [illegible]

Woolfolk Chemical Works, Inc.

Manufacturers **Ft. Valley, Ga.**

DIRECTIONS

[illegible] Do not apply or allow to drift to areas occupied by unprotected persons or beneficial animals. Workers entering treated areas the day of application should wear protective clothing.

[illegible] This product is toxic to fish, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of waste. Apply this product only as specified on this label.

[illegible] Dispose of waste pesticide and spillage by burying [illegible] and away from water supplies. Perforate or break empty con[illegible] with Agric[illegible]

[illegible] This product is toxic to bees and should not be applied when bees are [illegible] the area.

FIELD CROPS

TOBACCO: [illegible] - Budworm, Hornworm, Green Peach Aphid, Flea Beetle, Green Stink Bug and Cabbage Looper - apply at the rate of 1 to 2 quarts [illegible] for good coverage. For plant bed treatment, dilute [illegible] water and apply at the rate of 6 gallons per 100 [illegible] when insect infestation becomes damaging and [illegible] control. Do not apply within 5 days [illegible]

COTTON: [illegible] Weevil and Aphids - apply at the rate of 1 to 2½ [illegible] Budworm, Cabbage Looper, Cotton Stink Bug and [illegible] Begin applications when infestation re[illegible] as necessary for control. Do not [illegible] treated foliage to dairy or meat

VEGETABLES

BEANS: [illegible] Bean Beetle - apply at the rate of 1 quart per [illegible] when insects are first seen and repeat as necessary [illegible] than [illegible] times per season. Do not apply within 3