

769-275

08/06/1968

1/1

NET WEIGHT

SECURITY

4 1/2 POUNDS

50% THIODAN®

WETTABLE

ACTIVE INGREDIENT: *Endosulfan 50.0%
 INERT INGREDIENT: 50.0%

U. S. D. A. Reg. No. 769-275 TOTAL 100.0%
 *Hexachlorocycloheximethano - 2, 4, 3 - benzodioxathiepin oxide
 (R) Thiodan is a registered trademark of Farbwerke Hoechst A. C.
 U. S. Pat. No. 2,799,685

ACCEPTED

aug 6 1968

769-275
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC REASON REGISTERED
 UNDER NO. 769-275 SUBJECT
 TO ATTACHED DOCUMENTS.

WARNING! KEEP OUT OF REACH OF CHILDREN.

Hazardous if swallowed, inhaled or absorbed through skin. Do not breathe dust or spray mist. Do not get in eyes, on skin or on clothing. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking; wear clean clothing. During commercial or prolonged exposure in spray mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type passed by the United States Department of Agriculture for Thiodan protection. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Workers entering treated areas on the day of application should wear protective clothing.

PROTECT WILDLIFE: Toxic to fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on the label.

DIRECTIONS

FRUIT, NUT, AND FIELD CROPS

PEACHES: Peach Tree Borer and Lesser Peach Tree Borer - Use 1 1/2 pounds per 100 gallons of water and apply thoroughly to all bark area from ground to scaffold limbs. Apply as directed by local Agricultural Authorities. Usually two applications are required, one immediately after harvest and a second, 3 to 4 weeks later. Do not make more than 2 applications after shuck split. Do not apply within 21 days of harvest. Do not graze animals in treated orchards.

PECANS: Pecan Nut Casebearer and Black Pecan Aphid - Apply at 1 1/2 pounds per 100 gallons of water. For nut casebearer - apply when eggs of first generation appear on tips of young nuts which is usually about the time the tips of the small nuts have turned brown. A second application may be necessary after second generation eggs are deposited. Do not apply after shucks split. Do not graze meat or dairy animals in treated groves.

TOBACCO: Budworm, Hornworm, Green Peach Aphid, Flea Beetle, Green Stink Bug and Cabbage Looper - apply at the rate of 1 to 2 pounds per acre. Use sufficient water for good coverage. For plant bed treatment, use 1 pound to 100 gallons of water and apply at the rate of 6 gallons per 100 square yards. To control Green June Bug larvae in plant beds use 1/2 pound per 100 gallons of water and apply as a drench at first sign of larvae activity using 1 gallon per square yard of plant bed. Begin applications when insect infestation becomes damaging and repeat in 10 to 14 days or as necessary for control. Do not apply within 5 days of harvest.

VEGETABLES

APPLICATION: Make full coverage sprays using the recommended amount of Thiodan per acre in the amount of water necessary for good coverage.

BROCCOLI, CABBAGE: Cabbage Looper, Imported Cabbage Worm, Diamond-back Moth larvae and Cross-striped Cabbageworm - Use 1 1/2 to 2 pounds per acre. Flea Beetle, Harlequin Bug and Stink Bug - Use 1 1/2 pounds per acre. Do not apply within 7 days of harvest.

CUCUMBERS, MELONS, PUMPKINS, SQUASH: Aphids, Cucumber Beetles, Striped Cabbage Flea Beetle and Squash Vine Borer - Use 1 to 2 pounds per acre. For vine borer apply weekly to flower buds, stems and vines beginning when moths first appear. Do not apply within 1 day of harvest.

EGGPLANT, PEPPERS: Colorado Potato Beetle, Flea Beetle, Whitefly, Aphids, Green Stink Bug, Blister Beetle and Hornworms - Use 1 pound per acre. Apply when insects are first observed and repeat as necessary for control. Do not apply to eggplant or peppers within 1 day of harvest.

HEAD LETTUCE: Cabbage Looper, Diamond-back Moth larvae, Imported Cabbage Worm and Green Peach Aphid - Use 1 1/2 to 2 pounds per acre. Do not make more than 3 applications after thinning. Do not apply within 14 days of harvest. Do not feed crop refuse to livestock.

POTATOES, TOMATOES: Aphids, Colorado Potato Beetle, Flea Beetle, Green Stink Bug, Whitefly and Blister Beetles - Use 1 pound per acre. For Hornworm use 1 to 2 pounds per acre; for Tomato Russet Mite, Tomato Fruitworm, and Yellow Striped Armyworm - use 2 pounds per acre. Apply when insects first appear and repeat as required. Do not apply within 1 day of harvest. When applying in greenhouse, wear a mask or respirator of a type passed by the USDA for Thiodan protection.

DISPOSAL WARNING: Dispose of waste pesticide and container by burying in non-crop land away from water supplies. Never re-use container.

NOTICE

The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

WOOLFOLK CHEMICAL WORKS, LTD.

Manufacturers

860205

Fort Valley, Ga.