NET WEIGHT SECURITY 6 POUNDS

50% THIODAN WETTABLE

ACTIVE INGREDIENT: [illegible]
INERT INGREDIENT: [illegible]

[illegible] U.S.A. Reg. No. [illegible]
[illegible] is a registered trademark of Farbwerke Hoechst A. G.
U.S. Pat. No. 2,726,185

WARNING! KEEP OUT OF REACH OF CHILDREN.

Hazardous if swallowed, inhaled or absorbed through skin. Do not breathe dust or spray mist. Do not get in eyes, on skin or on clothing. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water. For eyes, get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking. Wear clean clothing. During commercial or prolonged exposure in spray mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type passed by the United States Department of Agriculture for Thiodan protection. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Workers entering treated areas on the day of application should wear protective clothing.

PROTECT WILDLIFE: Toxic to fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on the label.

PROTECT BEES: This product is toxic to bees and should not be applied when bees are actively visiting the area.

DIRECTIONS

FRUIT, NUT, AND FIELD CROPS

PEACHES: Peach Tree Borer and Lesser Peach Tree Borer - Use 1½ pounds per 100 gallons of water and apply thoroughly to all bark area from ground to scaffold limbs. Apply as directed by State Agricultural Extension Service. Usually two applications are required, one immediately after harvest and a second 3 to 4 weeks later. Do not make more than 2 applications after shuck split. Do not apply within 21 days of harvest. Do not graze animals in treated orchards.

PECANS: Pecan Nut Casebearer and Black Pecan Aphid - Apply at 1½ pounds per 100 gallons of water. For nut casebearer - apply when eggs of first generation appear on tips of young nuts which is usually about the time the tips of the small nuts have turned brown. A second application may be necessary after second generation eggs are deposited. Do not apply after shucks split. Do not graze meat or dairy animals in treated groves.

TOBACCO: Budworm, Hornworm, Green Peach Aphid, Flea Beetle, Green Stink Bug and Cabbage Looper - apply at the rate of 1 to 2 pounds per acre. Use sufficient water for good coverage. For plant bed treatment, use 1 pound to 100 gallons of water and apply at the rate of 4 gallons per 100 square yards. To control Green June Bug larvae in plant beds use 1 pound per 100 gallons of water and apply as a drench at first sign of larvae activity using 1 gallon per square yard of plant bed. Begin application when insect infestation becomes damaging and repeat in 10 to 14 days or as necessary for control. Do not apply within 5 days of harvest.

VEGETABLES

APPLICATION: [illegible] of the recommended rate per acre in enough water for adequate coverage.

BROCCOLI, CABBAGE: [illegible] Cabbage - Use 1½ pounds per acre. For control of Harlequin Bug, Green Stink Bug, Imported Cabbageworm and Cabbage Looper apply at 1½ pounds per acre. Do not apply within 7 days of harvest.

CUCUMBERS, MELONS, PUMPKINS, SQUASH: Aphids, Striped Cucumber Beetles, Spotted Cucumber Beetle, Flea Beetle and Squash Vine Borer - Use 1½ pounds per acre. For control of Aphids, Stink Bugs and Leaf Eating Beetles when first appear. Do not apply within [illegible] days of harvest.

EGGPLANT, PEPPERS: [illegible] Aphids, Green Stink Bug, Blister Beetle and Hornworms - Use 1½ pounds per acre. Apply when first appear. For control of Aphids, Stink Bug, Blister Beetle and Hornworms apply at 1½ pounds per acre. Do not apply within 7 days of harvest.

HEAD LETTUCE: Cutworms, Leaf Eating Beetles, Aphids, Harlequin Bug, Cabbage Looper and Armyworms - Use 1½ pounds per acre. For control of Cutworms, Leaf Eating Beetles, Aphids, Harlequin Bug, Cabbage Looper and Armyworms apply at 1½ pounds per acre. Do not apply within 14 days of harvest. Do not graze animals in treated fields.

POTATOES, TOMATOES: Aphids, Potato Leafhopper, Green Stink Bug, Blister Beetle, Potato Flea Beetles - Use 1½ pounds per acre. For control of Aphids, Potato Leafhopper, Green Stink Bug, Blister Beetle, Potato Flea Beetles and Yellow Striped Armyworm apply at 1½ pounds per acre. Apply when pest first appear. Do not apply within 3 days of harvest. When used on tomatoes should be applied at the rate of 1 pound per acre for the first 3 applications.

DISPOSAL WARNING: [illegible]

NOTICE

[illegible]

WOOLFOLK CHEMICAL WORKS, LTD.
Manufacturers 9nd115 **Fort Valley, Ga.**

169 375

APR 2-1 1971

U.S. C POISON FILE STEP
NO. SUBJECT
ATTACHED COMMENTS.