5% Sevin* - 75% Sulphur Dust

BEST AVAILABLE COPY

ACTIVE INGREDIENTS:	
Carbaryl (1-naphthyl N-Methylcarbamate)	5.0%
Sulphur, as elemental	75.0%
INERT INGREDIENTS:	20.0%
EPA REG. NO. 769-296 EPA EST. 769-GA-1	TOTAL 100.0%

* Trademark of Union Carbide Corp. for 1-naphthyl N-methylcarbamate.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention. **IF INHALED:** Remove victim to fresh air. **IF ON SKIN:** Wash with soap and water. Remove any contaminated clothing. **IF IN EYES:** Flush eyes with water. **NOTE TO PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-pam, epinine or cholinesterase inhibiting drugs.

PRECAUTIONARY STATEMENTS

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

APPLICATION: Begin applications when insects or damage is first observed and repeat on a regular 5 to 7 day schedule or as necessary for good insect control. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

BEANS: To control Mexican bean beetle, bean leaf beetle, cucumber beetles, leafhoppers, flea beetles, Japanese beetles, corn earworm and rust, apply at the rate of 20 pounds per acre. To control lygus bug, stink bug and Lima bean pod borer, apply 40 pounds per acre.

CARDOON, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI: To control flea beetles and Harlequin bug, apply at the rate of 20 pounds per acre. To control corn earworm, armyworm, cabbage caterpillars and small cabbage loopers, apply 20 to 40 pounds per acre.

OKRA: To control corn earworm and stink bugs, apply at the rate of 20 to 40 pounds per acre.

TOMATO, EGGPLANT, PEPPER: To control tomato fruitworm, fall armyworm, flea beetles and leafhoppers, apply at the rate of 20 pounds per acre. For control of tomato hornworm, European corn borer and stink bugs, apply at the rate of 20 to 40 pounds per acre.

POTATO: To control Colorado potato beetle, leafhoppers and flea beetles, apply at 20 pounds and for Lygus bug, apply at 40 pounds per acre.

INTERVAL BEFORE HARVEST: This product can be used up to and including the day of harvest on all vegetables listed except broccoli, Brussels sprouts, cabbage, cauliflower, kohlrabi, cowpea and okra. When applied to these vegetables, allow 3 days between the last application and harvest.

CAUTION: Do not apply this product to cucumbers, squash or melons because sulphur injury will result.

FORAGE CROPS

APPLICATION: Apply when diseases or insect damage appears and repeat 7 to 14 days later if necessary.

PEANUTS: To control Mexican bean beetle, potato beetles, thrips, leafhoppers, three-cornered alfalfa hoppers, cucumber beetles, bean leaf beetle, Japanese beetle, alfalfa caterpillar, velvet bean caterpillar and leaf spot, apply at the rate of 20 pounds per acre. To control armyworms, webworms, corn earworm, stink bugs, and grasshoppers, apply at the rate of 20 to 30 pounds per acre. Application may be made up to and including the day of harvest. Some leaf injury may occur if applications are made when tender foliage is wet.

ALFALFA, CLOVERS, SOYBEANS, COWPEAS: To control the same insects listed under peanuts, apply at the rate of 20 to 30 pounds per acre. Some leaf injury may occur if applications are made when tender foliage is wet.

Allow 3 days between the last application and harvest for alfalfa crops.

CORN (Forage and Fodder): To control corn earworm, European corn borer, fall armyworm, sap beetles and Japanese beetles, apply at the rate of 20 to 40 pounds per acre. Application may be made up to within 7 days of harvest. Application of this product to the tassel region of corn during pollen-shed period will seriously reduce bee populations.

SORGHUM (Grain and Forage): To control army worms, webworms, corn earworm, stink bugs and grasshoppers, apply at the rate of 20 to 40 pounds per acre. Allow 21 days between last application and harvest of grain, no time limit on sorghum forage.

STORAGE AND DISPOSAL

STORAGE: Store pesticide in original container and place in a locked storage area. Opened pesticide containers should be securely reclosed prior to storage. Protect from penetration of moisture, excessive heat or cold. Avoid cross-contamination with other pesticides or fertilizers. Use caution when storing pesticides in stacks.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE: SECURITY CHEMICAL COMPANY warrants that this product conforms to the chemical description on its label and is reasonably fit for the purposes stated on this label when used in accordance with the directions of the label and under normal conditions. SECURITY MAKES NO WARRANTIES OF MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, EXPRESSED OR IMPLIED, EXCEPT AS STATED ABOVE. Ineffectiveness or other unintended consequences may result because of such factors as environmental conditions, conditions of application and other conditions relating to the use and handling of this product. All such risks are assumed by the purchaser. SECURITY SHALL IN NO EVENT BE LIABLE FOR CONSEQUENTIAL OR INCIDENTAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NET WEIGHT: 50 LBS.

SECURITY CHEMICAL COMPANY

MANUFACTURERS ATLANTA, GEORGIA

ACCEPTED

JUN 28 1985

Under the Federal Insecticide, Fungicide and Rodenticide Act, as amended for the pesticide registered under EPA Reg. No. 769-296