SECURITY BRAND CHEMICALS · SECURITY BRAND CHEMICALS · SECURITY BRAND CHEMICALS

CONTROLS INSECTS AND DISEASES ON

SECURITY

DIRECTIONS

When this product is used as directed, it will control many of the important insects and diseases affecting fruits.

Use a good mechanical duster and apply to tree fruits at the rate of 5 to 8 ounces per tree, 1 pound per 100 feet of row to grapes and 3 ounces to 100 feet of row to strawberries. As a spray use 1¼ cups to 3 gallons of water and apply 1½ to 4 gallons per tree, 2½ gallons per 100 feet of row to grapes and 2 quarts per 100 feet of row on strawberries. Best results are obtained if the dust is applied in early morning or late afternoon when the air is calm. Do not apply during periods of high temperatures or high humidity. A one pound package is sufficient to make one application to three average size peach trees.

TIME OF APPLICATIONS

Begin applications at petal fall and repeat at 14 day intervals on peaches and plums and at 30 day intervals on cherries as necessary during the season. Do not apply to peaches within 14 days of harvest or to cherries and plums within 7 days of harvest.

APPLES AND PEARS: Begin applications 30 days after full bloom. If disease control is necessary earlier, apply separate sprays or dusts of captan or other suitable fungicide at petal-fall and two weeks later. Earlier applications may cause fruit thinning and on McIntosh and York varieties may cause some foliage injury. Repeat applications at 10 to 14 day intervals as necessary during the season. Do not apply to apples and pears within 7 days of harvest.

GRAPES: Make first application when new shoots are 2 to 4 inches long, the second application, when shoots are 8 inches long, the third, just before blossoms open, the fourth, just after blossoms fall, and additional applications at 10 to 14 day intervals as necessary during the season. Do not apply to grapes within 7 days of harvest.

STRAWBERRIES: Begin applications when plants start to bloom and repeat at 10 to 14 day intervals as necessary during the season. Do not apply within 2 days of harvest.

INSECTS AND DISEASES CONTROLLED

PEACHES, PLUMS AND CHERRIES: Plum curculio, Oriental fruit moth, Cat-facing bugs, Cherry fruit fly, Fruit tree leaf roller, Cherry aphid, Scale insects and mites, Brown rot, Peach scab, Rhizopus rot and Cherry leaf spot.

APPLES AND PEARS: Codling moth, Green apple aphid, White apple leafhopper, Apple mealybug, Apple maggot, Rosy apple aphid, Red banded leaf roller, Pear psylla, Periodical cicada, Eye spotted bud moth, European apple sawfly, Mites, Eastern tent caterpillar, Japanese beetle, Tentiform leaf miners, Forbes scale, San Jose scale, Oystershell scale and Lecanium scale, Scab, Brooks fruit spot, Sooty blotch, Fly speck, Bitter rot, Black rot, Botryosphaeria rot and Black pox.

GRAPES: Grape berry moth, Grape leaf folder, Grape leaf hopper and Japanese beetle, Black rot.

STRAWBERRIES: Meadow spittlebug and Strawberry leaf roller, Fruit rots.

VOLUME MEASUREMENTS FOR DILUTION

1 oz. by weight = 3 ½ Tablespoons
2 oz. by weight = ¼ Cup
6 oz. by weight = 1¼ Cups

NOTICE: The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

SECURITY BRAND CHEMICALS · SECURITY BRAND CHEMICALS · SECURITY BRAND CHEMICALS · SECURITY BRAND CHEMICALS · SECURITY BRAND CHEMICALS · SECURITY BRAND CHEMICALS

CONTROLS INSECTS AND DISEASES ON

SECURITY®

NET WEIGHT ONE POUND

PAN® FRUIT DUST OR SPRAY

CONTROLS INSECTS AND DISEASES ON PEACHES, PLUMS, APPLES, PEARS CHERRIES, GRAPES AND STRAWBERRIES

CONTAINS: *SEVIN • CAPTAN • *KELTHANE

ACTIVE INGREDIENTS

Carbaryl (1-naphthyl N-methylcarbamate)	7.50%
*Captan	7.50%
1,1-bis (p-chlorophenyl)-2,2,2-trichloroethanol	2.00%
INERT INGREDIENTS	83.00%
U.S.D.A. REG. NO. [illegible]	Total 100.00%

*N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide
U.S. Pat. Nos. 2,553,770, 2,553,771, 2,553,776
®Sevin is trademark of Union Carbide Corp. for 1-Naphthyl, N-Methylcarbamate
®Kelthane is trademark of Rohm & Haas

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Harmful if inhaled, swallowed, or absorbed through the skin. Do not breathe dust. Do not take internally. Avoid skin contact. Avoid getting in eyes. Wash hands before eating. Wash after contaminated clothing with soap and water before reuse. Avoid contamination of foodstuffs, feed troughs, and water receptacles. This product is toxic to honeybees. Avoid use during flowering and pollination period. Fish may be killed if their waters are contaminated with this product.

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidote.

WOOLFOLK CHEMICAL WORKS, LTD.
MANUFACTURERS • FORT VALLEY, GA.