

**ACCEPTED**  
 JUN 28 1985  
 Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act,  
 as amended, for the pesticide  
 registered under  
 EPA Reg. No. 769-304

Reg # 769-304  
 1123  
 1123

**PESTICIDE DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**NOTICE**

The directions for use given on this product are based upon use under normal and reasonably foreseeable conditions. If the directions for use are followed and the precautions set forth on the label are observed, research indicates that the product will give satisfactory results and that the residue will be within established tolerances. However, because of the fact that the climatic geographical and other conditions under which this product may be used will vary widely, the seller makes no warranty or representation, expressed or implied, concerning the material except that it conforms to the chemical description on the label and no person is authorized to make any other representation upon its behalf. The seller shall not be held responsible for damages resulting from the handling, storage or use of this product for any reason except the failure of the material to conform to the chemical description on the label. Damages, if any, which may result from the failure of this material to conform with the description on the label, shall be limited to direct damages, but shall not include consequential damages such as those resulting from crop or personal injury, residues in excess of official tolerances, or failure to give adequate control.

**NET WEIGHT: 50 POUNDS**

**SECURITY CHEMICAL COMPANY**  
**MANUFACTURERS ATLANTA, GEORGIA**



**Copper-Sulphur  
 Sevin®  
 Peanut Dust**

**For combined  
 Leaf Spot and Insect Control**

**ACTIVE INGREDIENTS:**

Sulphur, as elemental	80.0%
Copper, as metallic, from copper oxide	3.4%
carbaryl (1-naphthyl N-methylcarbamate)	5.0%
<b>INERT INGREDIENTS:</b>	<b>11.6%</b>
<b>TOTAL 100.0%</b>	

EPA REG. NO. 769-304      EPA EST. 769-GA-1  
 ® Trademark of Union Carbide Corp. for 1-naphthyl N-methylcarbamate

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION!**

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention. **IF INHALED:** Remove victim to fresh air. **IF ON SKIN:** Wash with soap and water. Remove any contaminated clothing. **IF IN EYES:** Flush eyes with water. **NOTE TO PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates or cholinesterase inhibiting drugs.

**PRECAUTIONARY STATEMENTS**

**CAUTION:** May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Subsequent plantings of food and feed crops in treated soil must be made with crops on this or other carbaryl labels. **PEANUTS:** To control thrips, leafhoppers, cucumber beetles, velvet bean caterpillar and leaf spot, apply at the rate of 20 pounds per acre. To control armyworms and corn earworms, apply at the rate of 20 to 30 pounds per acre. Apply when insects or signs of damage appear and repeat 7 to 14 days later if necessary. Application may be made up to and including the day of harvest. Some leaf injury may occur if applications are made when tender foliage is wet. For leaf spot control on Spanish peanuts, make the first application when spots appear on basal leaves. Usually 60 to 65 days after planting. Repeat application at two-week intervals until at least 3 applications have been made. On runner peanuts, apply the first dust about one week later than for Spanish then apply 3 additional applications at two-week intervals. If rain occurs within 24 hours of an application, repeat the application as soon as practical.

**STORAGE AND DISPOSAL**

**STORAGE:** Store pesticide in original container and place in a locked storage area. Opened pesticide containers should be securely resealed prior to storage. Protect from penetration of moisture, excessive heat or cold. Avoid cross-contamination with other pesticides or fertilizers. Use caution when storing pesticides in stacks.