


**CONTROLS**



**JAPANESE BEETLE**



**MEXICAN  
BEAN BEETLE**

**CONTROLS GARDEN and ORNAMENTAL  
INSECT PESTS**

**NET WT. 4 LBS.**

<b>ACTIVE INGREDIENT:</b>	
Carbaryl (1-naphthyl N-methylcarbamate) .....	10.0%
<b>INERT INGREDIENTS:</b> .....	<u>90.0%</u>
<b>TOTAL</b>	<b>100.0%</b>

EPA Reg. No. 769-315



**CORN:** To control Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetles, and Japanese Beetles - apply at the rate of 15 to 20 pounds per acre. For European Corn Bo. apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until tassels emerge. For Budworm damage in the whorl, apply to entire plant and repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending upon severity of infestation. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

Continued from Back Panel

#### EARWIGS

This material is effective in controlling earwigs and when properly used will reduce the number gaining entrance to homes and buildings. Apply heavily at the rate of 1 pound to 150 linear feet completely around the foundation wall of buildings. The material should be applied heavily in a band 3 to 4 inches in width, using a shaker type can or fruit jar with holes punched in the lid. Earwigs breed over wide areas around yards and fields and are driven by cold weather to seek shelter in warm buildings. As they move along the foundation wall seeking entrance, they come in contact with the treated barrier and are killed. Be sure to treat in front of all steps and other entrance ways as well as along the foundation wall. Treat, also, around refuse piles, lumber mulch, and other areas in the yard offering areas of concealment to earwigs.

Repeat application in 10 days or when earwigs become troublesome. If the dust barrier is washed away by rain, it should be replaced with a new application if earwigs are still a problem.

**KEEP OUT OF REACH OF CHILDREN.** Harmful if inhaled, swallowed, or absorbed through the skin. Do not breathe dust. Do not take internally. Avoid skin contact. Avoid getting in eyes. Wash hands before eating. Wash all contaminated clothing with soap and water before re-use. Avoid contamination of foodstuffs, feed, and water receptacles.

**NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. See package insert for details.

**PROTECT BEES:** This material is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

## DIRECTIONS

### VEGETABLES

**APPLICATION:** Begin applications when the first damage is first observed and repeat on a regular 5 to 7 day schedule or as necessary for good insect control. In gardens apply at the rate of 1 ounce to 50 feet of row, which is equivalent to 20 pounds per acre.

**BEANS:** To control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetles, Leafhoppers, Flea Beetles, Japanese Beetles, and Corn Earworm—apply at the rate of 10 pounds per acre. To control Lygus Bug, Stink Bug and Lima Bean Pod Borer—apply 20 pounds per acre.

**CABBAGE, COLLARDS, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, COLESLAW, MUSTARD GREENS, TURNIPS, KOHLRABI, and SPINACH:** To control Flea Beetle and Harlequin Bug—apply at the rate of 10 pounds per acre. To control Corn Earworm, Armyworm, Imported Cabbageworm, Diamond-back moth larvae—apply 10 to 20 pounds per acre.

**CUCUMBERS, SQUASH, CANTALOUPE, PEPPERS, MELONS:** To control Pickleworm, Cucumber Borer, Cucumber Beetles, Squash Bug, Flea Beetles—apply at the rate of 10 pounds per acre.

**CORN:** To control Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetles, and Japanese Beetles—apply at the rate of 15 to 20 pounds per acre. For Budworm damage in the whorl, apply to entire plant and repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending upon severity of infestation.

**CUCUMBERS:** To control Flea Beetles, Leafhoppers—apply at the rate of 10 to 20 pounds per acre.

**LETTUCE:** To control Flea Beetles, Harlequin Bug, and Leafhoppers—apply at the rate of 10 pounds per acre. To control Corn Earworm, Armyworm, Imported Cabbageworm, Diamond-back moth larvae, Lygus Bugs—apply at the rate of 10 to 20 pounds per acre.

**POTATO:** To control Corn Earworm and Stink Bugs—apply at the rate of 10 to 20 pounds per acre.

**VEGETABLES:** To control European Corn Borer, Fall Armyworm, Flea Beetles, Japanese Beetles—apply at the rate of 10 pounds per acre. For control of Tomato Hornworm, European Corn Borer and Stink Bugs—apply at the rate of 10 to 20 pounds per acre.

**POTATO:** To control Colorado Potato Beetle, Leafhoppers, and Flea Beetles, apply at 10 pounds and for Lygus Bug, apply at 20 pounds per acre.

**INTERVAL BEFORE HARVEST:** Allow 3 days between application and harvest on Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kohlrabi and Head Lettuce. Allow 14 days on Collards, Leaf Lettuce, Mustard Greens, Spinach and Turnips. On other vegetables listed on this label application can be made up to and including the day of harvest.

### ORNAMENTALS

To control Apple Aphid, Bagworm, Birch Leaf Miner, Gopher Beetles, Boxelder Bug, Boxwood Leaf Miner, Elm Leaf Aphid, Elm Leaf Beetle, Flea Beetles, Japanese Beetle, Lace Bugs, Leafhoppers, Leaf Rollers, Oak Leaf Miners, Orange Tortrix, Periodical Cicada, Plant Bugs, Psyllids, Rose slug, Rose Aphid, Scale insects in the crawler stage, Tent Caterpillars, Willow Leaf Beetle and exposed Thrips—apply thoroughly to the infested plants at first sign of damage and repeat at 7-day intervals or as necessary for good insect control. To obtain scale control apply in Spring and early Summer when scale crawlers are present.

### LAWNS

To control Leafhoppers, Fall Armyworm, and Sod Webworm—apply at the rate of 2 pounds per 1,000 square feet. For best results, apply after rain or watering and do not water for at least 2 days after application. Repeat in 2 to 3 weeks if necessary.

**NOTICE:** The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

**DISPOSAL WARNING:** Dispose of waste pesticide and container in a safe place. Never re-use container.

**CAUTION:** Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application.

Continued on Side Panel