

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

TOP: READ THE LABEL



CAN KILL YOU

DANGER!
Keep out of reach of children

POISON
See side panels for antidote and precautions

ACU...
769-329
JUL 7 1982

THREE WAY COTTON SPRAY 6-3

ACTIVE INGREDIENTS:

- Methyl Parathion (Dichloro) 1.00%
- Resmethion 1.00%
- Ethion Parathion (Dichloro) 1.00%
- Resmethion 1.00%
- Resmethion 1.00%

INERT INGREDIENTS:

- EPA Reg. No. 100-100-01
- Trade Name: ...
- Product: ...

NOT FOR HOME USE

DANGER-POISON-PRECAUTIONS



CAN KILL YOU BY SKIN CONTACT

This product can kill you if touched by hands or sprayer or splashed on skin, or eyes or clothing. Liquid goes through clothes.



CAN KILL YOU IF BREATHED

This product can kill you if you breathe it in.



CAN KILL YOU IF SWALLOWED

This product can kill you if you swallow it.

WOOLFOLK CHEMICAL WORKS INC
286250

Fort Valley GA 31030

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
STATEMENT OF PRACTICAL TREATMENT**

... IF SWALLOWED: ... IF INHALED: ... IF ON SKIN: ... IF IN EYES: ...

POISON SIGNS (symptoms)

Parathion is a very dangerous poison. It rapidly enters the body of contact with skin surfaces and eyes. Clothing wet with this material must be removed immediately. Exposed persons must receive prompt medical treatment if they may do.

Some of the symptoms of poisoning are: headache, nausea, vomiting, Cramps, Weakness, Blurred Vision, Pinpoint Pupils, Tightness of Chest, Labored Breathing, Nervousness, Sweating, Watery Eyes, Tremors or Flushing of Mouth and Nose, Muscle Spasms and Convulsions.

NOTE TO PHYSICIAN

Antidote: atropine sulfate (1 mg per dose). Two to FOUR mg intravenously or orally (adults) as a first dose. Repeat every 5 to 10 minutes until atropine is administered in 2 PAM or 4 mg (children) and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion causes a decrease in blood clotting time and peripheral vascular constriction and produces a fall in blood pressure. Atropine should be given early, before the patient reaches the stage of respiratory paralysis and treated with artificial respiration and about 10 mg of atropine. Atropine should be given in 1 mg increments until the patient is breathing VERY CLOSELY. REPEAT SIGNS OF THE PATIENT INDICATES REPEAT AT LEAST 48 HOURS.

USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE EQUIPMENT AND CLOTHING

... (Detailed list of protective equipment and clothing requirements)


WORK SAFETY RULES

... (Detailed list of work safety rules, including instructions on handling, storage, and disposal)

ENVIRONMENTAL HAZARDS

... (Information regarding environmental hazards and disposal procedures)

**PARATHION
MIXTURE,
LIQUID
NA2783 RQ**



POST TREATED AREAS
DIRECTIONS FOR USE
COTTON

[Faint, mostly illegible text follows, likely containing detailed application instructions for cotton.]

STORAGE & DISPOSAL
NOTICE
**DISCLAIMER OF WARRANTY
MERCHANTABILITY AND LIABILITY**

[Faint text follows, detailing storage, disposal, and liability information.]

769-329