

ACCEPTED

MAY 12 1976

UNDER THE FEDERAL PESTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTRATION  
NUMBER NO. 769-336

SECURITY

# 20% CHLORDANE

GRANULATED

## KILLS FIRE ANTS

**ACTIVE INGREDIENT:**

\*Technical Chlordane ..... 20.0%

**INERT INGREDIENTS:** ..... 80.0%

EPA Reg. No. 769-336      EPA Est. 769-GA-1      TOTAL 100.0%

\*Equivalent to 12% Octachloro-4, 7-methanotetrahydroindane and 8% related compounds.

**KEEP OUT OF REACH OF CHILDREN.**

**CAUTION!**

SEE ADDITIONAL PRECAUTIONARY STATEMENTS BELOW.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

KEEP OUT OF REACH OF CHILDREN. Avoid inhaling dust. Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid contamination of feed and foodstuffs. If spilled on skin, wash thoroughly with soap and water. Do not allow livestock to graze treated areas. Do not feed treated forage to dairy animals or animals being finished for slaughter. Keep children and pets off treated areas until this material is washed into the soil and the grass is dry.

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of ponds, lakes, or streams. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**FIRE ANTS**

To control fire ants in noncropland only (lawns, ornamental turf, cemeteries, golf courses, parks, industrial sites, roadsides and similar noncropland).

FOR BROADCAST SOIL TREATMENT apply 5 to 7½ lbs. (1 to 1½ lbs. technical chlordane) per acre. Make broadcast application. Cover surface uniformly.

Do not treat soil when it is wet. Do not apply to pastures or rangeland. Do not contaminate feed and food crops.

FOR MOUND APPLICATION apply 4/5 to 1-2/3 cups Chlordane 20% (equivalent to 0.07 to 0.14 lbs. technical chlordane) to entrance of nest as well as to an area 3 feet around mound. Treated areas should be examined 3 weeks after treatment and surviving colonies retreated. Do not graze livestock on treated areas within 1 year of treatment application. Do not contaminate feed and food crops.

**WHITE-FRINGED BEETLE LARVAE**

(Limited to the 8 Southeastern states of Alabama, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina, and Tennessee.)

BROADCAST TREATMENT - Cropland to be planted to beans, beets, blackeyed peas, cabbage, cantaloupe, collards, cowpeas, cucumbers, eggplant, peppers, rutabaga, sweet potatoes, turnips, kale, lettuce, melons, mustard greens, okra, onions, peas, pimientos, radishes, spinach, squash, tomatoes, and watermelons can be treated prior to planting. Apply to soil surface at the rate of 1¼ lbs. per 1000 square feet (50 lbs. per acre) and work into the upper 2 to 3 inches of soil.

**JAPANESE BEETLE LARVAE**

(Limited to use under Federal/State Quarantine Programs.)

NURSERY STOCK - Apply to soil surface at the rate of 1¼ lbs. per 1000 square feet.

STORAGE - Do not store near feed or foodstuffs.

**STORAGE** - Do not store near feed or foodstuffs. Keep out of reach of children.

**DISPOSAL** - Dispose of waste pesticides and container in a safe place away from water supplies. Never re-use container.

**WARRANTY STATEMENT**

If for any reason this product is not used, applied, stored or disposed of in accordance with instructions set forth on this label, then and thereafter Woolfolk Chemical Works, Inc. terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness.

**Manufacturers**

**WOOLFOLK CHEMICAL WORKS, INC.**

**Ft. Valley, Ga.**