

NET WEIGHT



50 POUNDS

# 6% MANEB DUST

ACCEPTED

May 8, 69

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 769-342

ACTIVE INGREDIENT:

Manganese ethylenebisdithiocarbamate (Maneb) ..... 6.0%

**INERT INGREDIENTS:** ..... 94.0%

U. S. D. A. Reg. No. 769-342

TOTAL 100.0%

FOR USE ON LIMA BEANS, SNAP BEANS, TOMATOES, EGGPLANT,  
PEPPERS, CUCUMBERS, CANTALOUPE, MELONS, SQUASH, WATERMELONS

## CAUTION! KEEP OUT OF REACH OF CHILDREN.

May cause irritation of nose, throat, eyes and skin. Do not breathe dust. In case of skin contact, wash with water; for eyes, get medical attention.

**PROTECT WILDLIFE:** This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label.

**STORAGE:** Store in cool, dry place. Do not allow to become wet or overheated in storage. This may cause chemical changes which will impair the fungicidal effectiveness of the product.

## DIRECTIONS

**LIMA BEANS, SNAP BEANS, DRY BEANS:** To control Downy Mildew, Anthracnose and Rust, apply at the rate of 30 pounds per acre. Begin applications when 5 to 10% of blooms show and continue at 7-day intervals as necessary. Do not apply to snap beans (String beans) within 4 days of harvest. For early season insect and disease control, use Maneb-Parathion-DDT Vegetable Dust and for combined insect and disease control anytime, use Vege-O-Gard.

**TOMATOES:** To control Early Blight, Late Blight, Anthracnose, Septoria Leaf Spot, and Gray Leaf Spot (Stemphylium), apply at the rate of 30 to 40 pounds per acre. Begin applications when first cluster fruit is well formed and repeat at 7 to 10-day intervals.

Earlier applications and more frequent applications may be necessary if disease conditions are unusually severe. For early season insect and disease control, use Maneb-Parathion-DDT Vegetable Dust and for combined insect and disease control anytime, use Vege-O-Gard.

**EGGPLANT, PEPPERS:** To control Fruit Rot and Anthracnose, apply at the rate of 20 to 25 pounds per acre. Begin when fruit begins to form or earlier if disease conditions are unusually severe. For early season insect and disease control, use Maneb-Parathion-DDT Vegetable Dust and for combined insect and disease control anytime, use Vege-O-Gard.

**CUCUMBERS, CANTALOUPE, MELONS, SQUASH, WATERMELONS:** To control Downy Mildew, Anthracnose and Gummy stem blight, apply at the rate of 15 to 50 pounds per acre. Start applications when runners begin to form or at the first sign of disease. Repeat at weekly intervals. Direct dust to underside of leaves.

### DISPOSAL WARNING

Dispose of waste pesticide and container by burying in non-crop land away from water supplies. Never re-use container.

## NOTICE

### DISCLAIMER OF WARRANTY, MERCHANTABILITY AND LIABILITY

If for any reason this product is not used, applied, stored, transported or disposed of in accordance with the instructions set forth on this label, then and thereafter Woolfolk Chemical Works, Ltd. terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness.

# WOOLFOLK CHEMICAL WORKS, LTD.

MANUFACTURERS

960415

FORT VALLEY, GEORGIA