

SECURITY BRAND  
6% MANEB DUST

FOR USE ON LIMA BEANS, SNAP BEANS, DRY BEANS, TOMATOES, EGGPLANT, PEPPERS, CUCUMBERS,  
CANTALOUPE, MELONS, SQUASH AND WATERMELONS

ACTIVE INGREDIENT:

Manganese ethylenebisdithiocarbamate (Maneb) ----- 6%

INERT INGREDIENTS: ----- 94%

TOTAL 100%

EPA Reg. No. 769-342

EPA Est. 769-GA-1

KEEP OUT OF REACH OF CHILDREN  
CAUTION

HAZARDS TO HUMANS & DOMESTIC ANIMALS

May cause irritation of eyes, nose, throat and skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. In case of contact, flush with plenty of water. For eyes, get medical attention.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Induce vomiting by immediately giving two glasses of water and sticking finger in throat. Call a physician immediately. IN INHALED: Move victim to fresh air. IF ON SKIN: Wash skin thoroughly with soap and water. IF IN EYES: Flush eyes with plenty of water and get prompt medical attention.

ENVIRONMENTAL HAZARDS

Toxic to fish. Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

NOTICE: When mixing or loading this product, all workers must wear the following protective clothing: long pants, long-sleeved shirt, gloves, hat and boots.

VEGETABLES

LIMA BEANS, SNAP BEANS, DRY BEANS: To control downy mildew, anthracnose and rust, dust at the rate of 30 pounds per acre. Begin applications when 5 to 10% of blooms show and continue at 7-day intervals as necessary. Do not apply to snap beans (string beans) within 4 days of harvest.

TOMATOES: To control early blight, late blight, anthracnose, septoria leaf spot, and gray leaf spot (stemphylium), dust at the rate of 30 to 40 pounds per acre. Begin applications when first cluster fruit is well formed and repeat at 7 to 10-day intervals. Earlier applications and more frequent applications may be necessary if disease conditions are unusually severe.

EGGPLANT, PEPPERS: To control fruit rot and anthracnose, dust at the rate of 20 to 25 pounds per acre. Begin when fruit begins to form or earlier if disease conditions require it, and continue at weekly intervals as necessary.

CUCUMBERS, CANTALOUPE, MELONS, SQUASH, WATERMELONS: To control downy mildew, anthracnose and gummy stem blight, dust at the rate of 15 to 50 pounds per acre. Start applications when runners begin to form or at the first sign of disease. Repeat on weekly intervals. Direct dust to underside of leaves.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. PESTICIDE DISPOSAL: Pesticide that cannot be used according to label instructions must be disposed of according to federal, state or local procedures. CONTAINER DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bag in a sanitary landfill or by incineration if allowed by state and local authorities.

NOTICE: The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

WOOLFOLK CHEMICAL WORKS, INC.

MANUFACTURERS

FORT VALLEY, GEORGIA