

769-345 TM 12  
131

the day of harvest. Some leaf injury may occur if applications are made when tender foliage is wet. For leaf spot control on Spanish peanuts, make the first application when spots appear on basal leaves, usually 60 to 65 days after planting. Repeat application at two-week intervals, until at least three applications have been made. On runner peanuts, apply the first dust about one week later than for Spanish; then apply three additional applications at two-week intervals. If rain occurs within 24 hours of an application, repeat the application as soon as practical.

**STORAGE AND DISPOSAL**

**STORAGE:** Store pesticide in original container and place in a locked storage area. Opened pesticide containers should be securely resealed prior to storage. Protect from penetration of moisture, excessive heat or cold. Avoid cross-contamination with other pesticides or fertilizers. Use caution when storing pesticides in stacks.

**PESTICIDE DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment then dispose of empty bag in a sanitary landfill or by incineration of, if allowed by state and local authorities, by burning. If burned, stay out of smoke


**NOTICE**

**DISCLAIMER OF WARRANTY  
MERCHANTABILITY AND LIABILITY**

If for any reason this product is not used, applied, stored, transported or disposed of in accordance with the instruction set forth on this label, then and thereafter Security Chemical Company terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness.

**\*NET WEIGHT: 50 POUNDS\***

**SECURITY CHEMICAL COMPANY  
MANUFACTURERS ••••• ATLANTA, GEORGIA**



**Copper - Sulphur  
10% Sevin®  
Peanut Dust**

**For combined Leaf Spot  
and Insect Control.**

**ACTIVE INGREDIENTS:**

- Sulphur, as elemental ..... 72.5%
- Copper, as metallic, from copper oxide ..... 3.4%
- Carbaryl (1-naphthyl N-methylcarbamate) ..... 10.0%

**INERT INGREDIENTS:** ..... 14.1%


EPA Reg. No 769-345

EPA Est. 769-GA-1

® Trademark of Union Carbide Corp. for 1-naphthyl N-methylcarbamate.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

(See additional precautionary statements on side panel.)



**IF SWALLOWED:** Drink 1 to 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

**IF INHALED:** Remove victim to fresh air.

**IF ON SKIN:** Wash with soap and water. Remove any contaminated clothing.

**IF IN EYES:** Flush eyes with water.

**NOTE TO PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates or cholinesterase-inhibiting drugs.

**PRECAUTIONARY STATEMENTS**

**CAUTION:** May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of waste. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

**PEANUTS:** To control Mexican bean beetle, blister beetles, thrips, leafhoppers, three-cornered alfalfa caterpillar, velvet bean caterpillar and leaf spot, apply at the rate of 10 pounds per acre. To control armyworms, webworms, corn earworms, stink bugs and grasshoppers, apply at the rate of 10 to 15 pounds per acre. Apply when insects or signs of damage appear and repeat 7 to 14 days later if necessary. Application may be made up to and including