

Jun 15, 71

SECURITY

10% SEVIN® - 75% SULPHUR DUST

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-Methylcarbamate) 10.0%
Sulphur, as elemental 75.0%

INERT INGREDIENTS: 15.0%

U. S. D. A. Reg. No. 769-349 TOTAL 100.0%
(R) Trademark of Union Carbide Corp. for 1-naphthyl N-Methylcarbamate.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with eyes or skin. Wash thoroughly after handling. Wash all contaminated clothing with soap and water before reuse. Do not store near feed and foodstuffs. Sulphur may burn foliage when temperatures are excessively high.

ADDITIONAL PHYSICIAN: Carbaryl is a moderate to strong cholinesterase inhibitor. Atropine is antidotal.

AVOID CONTAMINATION OF FOODSTUFFS: Avoid contamination of feed, feeding troughs, and watering receptacles.

INSECTICIDE RESIDUES: This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS

VEGETABLES

APPLICATION: Begin applications when insect or damage is first observed and repeat on a regular 5 to 7 day schedule or as necessary for good insect control.

BEANS: To control Mexican Bean Beetle, Bean Leaf Bee,

tle, Cucumber Beetles, Leafhoppers, Flea Beetles, Japanese Beetles, Corn Earworm and Rust - apply at the rate of 10 pounds per acre. To control Lygus Bug, Stink Bug and Lima Bean Pod Borer - apply 20 pounds per acre.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI: To control Flea Beetle and Harlequin Bug - apply at the rate of 10 pounds per acre. To control Corn Earworm, Armyworm, Cabbage Caterpillars and small Cabbage Loopers - apply 10 to 20 pounds per acre.

OKRA: To control Corn Earworm and Stink Bugs - apply at the rate of 10 to 20 pounds per acre.

TOMATO, EGGPLANT, PEPPER: To control Tomato Fruitworm, Fall Armyworm, Flea Beetles and Leafhoppers - apply at the rate of 10 pounds per acre. For control of Tomato Hornworm, European Corn Borer and Stink Bugs - apply at the rate of 10 to 20 pounds per acre.

POTATO: To control Colorado Potato Beetle, Leafhoppers, and Flea Beetles, apply at 10 pounds and for Lygus Bug, apply at 20 pounds per acre.

INTERVAL BEFORE HARVEST: This product can be used up to and including the day of harvest on all vegetables listed except Broccoli, Brussels Sprouts, Cabbage, Cauliflower, and Kohlrabi. When applied to these vegetables allow 7 days between the last application and harvest.

CAUTION: Do not apply this product to Cucumbers, Squash or Melons because sulphur injury will result.

FORAGE CROPS

APPLICATION: Apply when disease or insect damage appears and repeat 7 to 14 days later if necessary.

PEANUTS: To control Mexican Bean Beetle, Blister Bee-

ties, Thrips, Leafhoppers, Three-Cornered Alfalfa Hopper, Cucumber Beetles, Bean Leaf Beetle, Japanese Beetle, Alfalfa Caterpillar, Velvet Bean Caterpillar and Leaf Spot - apply at the rate of 10 pounds per acre. To control Armyworms, Webworms, Corn Earworm, Stink Bugs, and Grasshoppers, apply at the rate of 10 to 15 pounds per acre. Application may be made up to and including the day of harvest. Some leaf injury may occur if applications are made when tender foliage is wet.

ALFALFA CLOVER AND COWPEAS: To control the insects listed under forage crops apply at the rate of 10 to 15 pounds per acre. Some leaf injury may occur if applications are made when tender foliage is wet.

CORN: (Forage and Fodder) To control Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetles and Japanese Beetles, apply at the rate of 10 to 20 pounds per acre. Application may be made up to within 7 days of harvest. Application of this product to the tassel region of corn during pollen shed period will seriously reduce bee populations.

SORGHUM: (Grain and Forage) To control Armyworms, Webworms, Corn Earworm, Stink Bugs, and Grasshoppers, apply at the rate of 10 to 20 pounds per acre. Allow 21 days between last application and harvest of grain, no time limit on sorghum forage.

Dispose of waste pesticide and container by burying in non-crop land away from water supplies. Never re-use container.

NOTICE

The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

Manufacturers

WOOLFOLK CHEMICAL WORKS, LTD.

Fort Valley, Ga.