

NET WEIGHT



50 POUNDS

6% BOTRAN® PEACH DUST

ALSO FOR CONTROL OF CERTAIN DISEASES OF GRAPE, STRAWBERRY,
TOMATO, ONION, GARLIC, AND ORNAMENTALS

aug 8 1967
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROENTICIDE ACT.
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 769-355 SUBJECT
TO REVISION COMMENTS.

ACTIVE INGREDIENT: 2, 6-dichloro-4-nitroaniline 6.0%
INERT INGREDIENTS: 94.0%

U. S. D. A. Reg. No. 769-355
(R) Trademark Tuco Products Co., Div. of the Upjohn Co.,
Kalamazoo, Michigan

TOTAL 100.0%

WARNING: KEEP OUT OF REACH OF CHILDREN.
Harmful if swallowed. Avoid inhalation of dust. Avoid contact with skin, eyes and clothing. In case of contact, remove contaminated clothing and wash skin with soap and water. Avoid contamination of feed and foodstuffs.

DIRECTIONS

CAUTION: Do not apply when weather conditions favor drift of dust from crop area treated. Keep out of any body of water. Fish may be killed.

PEACHES: To control Brown Rot and Rhizopus Rot apply at the rate of 25 to 40 pounds per acre. Make 3 applications, the first 18 days before harvest, the second 10 days before harvest and the third 1 day before harvest. Do not apply closer than 1 day of harvest.

GRAPE: To control Botrytis rot apply at the rate of 30 pounds per acre. Begin applications while clusters are still loose and repeat applications at two-week intervals. Allow one day between last application and harvest.

STRAWBERRY: To control Botrytis rot and Rhizopus rot apply at the rate of 25 pounds per acre. Begin application when new leaves are expanded and blossom buds are visible. Repeat applications at weekly intervals through harvest.

TOMATO: In the greenhouse to control Stem Canker (Botrytis) apply at the rate of 30

pounds per acre. Dust stem of plant from ground level up to a height of 18 to 24 inches. Repeat weekly or as necessary.

ONION and GARLIC: To control White Rot (Sclerotium) apply at the rate of 250 to 400 pounds per acre. For direct seeding incorporate the dust in top 1½ inches of soil before onions are seeded. When planting sets or buds dust into furrow so treated soil surrounds sets or buds.

ORNAMENTALS

ROSE, HYDRANGEA: To control Botrytis Blight apply in the field at the rate of 30 pounds per acre prior to shipment or storage. For storage apply thoroughly to storage area and to all canes and dead foliage; repeat application during long storage periods.

CHRYSANTHEMUM, GERANIUM, ROSE: To control Botrytis Blight apply at the rate of 30 pounds per acre to stock cuttings or greenhouse plants. Begin applications when disease is anticipated or first appears. Dust foliage and flowers at 7 to 14-day intervals. During prolonged damp, cool periods, apply every 5 to 7 days.

GLADIOLUS: To control Dry Rot (Stromatinia) dust at the rate of 250 pounds per acre into furrows at planting time so treated soil surrounds corms. To control Botrytis Blight apply at the rate of 30 pounds per acre. Force dust between spikes. Apply to bunched cut flower spikes after harvest and before shipment or storage.

NOTICE

Woolfolk Chemical Works, Ltd. makes no warranty, express or implied, concerning this material, except that it conforms to the chemical description on the label. Neither Woolfolk Chemical Works, Ltd. nor the seller shall be held responsible in any manner for any personal injury or property damage or other type of loss resulting from the handling, storage or use of this material. The buyer assumes all risk and liability therefrom and accepts and uses this material on these conditions.

WOOLFOLK CHEMICAL WORKS, LTD.

MANUFACTURERS

660512

FORT VALLEY, GEORGIA