

SECURITY

6% BOTRAN DUST

ACTIVE INGREDIENT

2,6-DIBROMO-4-NITROANILINE

INERT INGREDIENTS

EPA Reg. No. 769-356

60

64.9

TOTAL

100.0

BOTRAN is the registered trademark of Woolfolk Chemical Works, Inc., Fort Valley, Georgia.

CONTROLS CERTAIN DISEASES ON PEACHES, GRAPES, TOMATOES, ONIONS, GARLIC AND ORNAMENTALS

KEEP OUT OF REACH OF CHILDREN
CAUTION

In case of eye contact, flush eyes with plenty of water. Get medical attention if irritation persists.

HAZARD TO HUMANS & DOMESTIC ANIMALS
CAUTION

Avoid contact with skin, eyes, nose, mouth. Wash thoroughly after handling.

ENVIRONMENTAL HAZARD

This pesticide is toxic to fish. Do not apply directly to water, to areas that drain to water, or to areas adjacent to water. Do not apply within 100 feet of any water body. Do not apply within 100 feet of any water body.

DIRECTIONS FOR USE
GENERAL CLASSIFICATION

This is a complete list of crops and diseases for which this pesticide is registered.

METHOD OF APPLICATION Mix with water and apply as a dust. Do not apply when weather conditions favor drift.

PEACHES To control Brown Rot, apply at the rate of 25 to 30 pounds per acre at 14 to 16 weeks before harvest. Do not apply more than once per season. Do not apply within 14 days before harvest. Do not apply to fruit that is being harvested. Do not apply to fruit that is being harvested.

GRAPES To control Botrytis Rot, spray at the rate of 50 pounds per acre at 14 to 16 weeks before harvest. Do not apply more than once per season. Do not apply to fruit that is being harvested.

TOMATOES To control Stem Canker, Botrytis, apply at the rate of 25 to 30 pounds per acre at 14 to 16 weeks before harvest. Do not apply more than once per season. Do not apply to fruit that is being harvested.

ONION and GARLIC To control Botrytis, apply at the rate of 250 to 300 pounds per acre at 14 to 16 weeks before harvest. Do not apply more than once per season. Do not apply to fruit that is being harvested.

ORNAMENTALS

ROSE, HYDRANGEA To control Botrytis, apply in the field at the rate of 25 to 30 pounds per acre at 14 to 16 weeks before harvest. Do not apply more than once per season. Do not apply to fruit that is being harvested.

CHLAYSANTHEMUM, GERANIUM, ROSE To control Botrytis, apply in the field at the rate of 25 to 30 pounds per acre at 14 to 16 weeks before harvest. Do not apply more than once per season. Do not apply to fruit that is being harvested.

GLADIOLUS To control Botrytis, apply in the field at the rate of 250 pounds per acre at 14 to 16 weeks before harvest. Do not apply more than once per season. Do not apply to fruit that is being harvested.

STORAGE & DISPOSAL

Store in original container. Do not use other containers. Do not use if container is damaged. Do not use if contents are empty. Do not use if contents are solid.

NOTICE This pesticide is registered for use on the crops and diseases listed above. Do not use on other crops or diseases.

WOOLFOLK CHEMICAL WORKS, INC.
MANUFACTURERS **BEST DOCUMENT AVAILABLE** **FORT VALLEY, GA**

SECURITY

6% BOTRAN DUST

Net Weight 50.00
Gross Weight 54.00
EPA Est. 769 GA 1 TOTAL 100.0

Trademark of TUCD Products Co., Div. of the Upjohn Co., Kalamazoo, Michigan

RETAIN DISEASES ON PEACHES, GRAPES, TOMATOES, ONIONS, GARLIC AND ORNAMENTALS

KEEP OUT OF REACH OF CHILDREN CAUTION

Flush eyes immediately with plenty of water. Get medical attention.

HAZARD TO HUMANS & DOMESTIC ANIMALS
CAUTION

with skin, eyes or clothing. Wash thoroughly after treatment.
ENVIRONMENTAL HAZARD
Do not use with care when applying in areas adjacent to any body of water and streams. Do not apply when weather conditions favor drift or runoff.

GRAPES To control botrytis Rot, apply Botran at the rate of 2 1/2 to 3 1/2 lbs. per acre of fruit. Do not apply when flowers are present. Do not apply to fruit that is to be eaten. Allow one day between last application and harvest.

TOMATOES To control blight, apply Botran at the rate of 2 1/2 to 3 1/2 lbs. per acre. Do not apply to fruit that is to be eaten. Allow one day between last application and harvest.

ONION and GARLIC To control White Rot, apply Botran at the rate of 2 1/2 to 3 1/2 lbs. per acre. Do not apply to fruit that is to be eaten. Allow one day between last application and harvest.

ORNAMENTALS

ROSE, HYDRANGEA To control Botrytis, apply Botran at the rate of 2 1/2 to 3 1/2 lbs. per acre. Do not apply to fruit that is to be eaten. Allow one day between last application and harvest.

CHRYSANTHEMUM, GERANIUM, ROSE To control Botrytis, apply Botran at the rate of 2 1/2 to 3 1/2 lbs. per acre. Do not apply to fruit that is to be eaten. Allow one day between last application and harvest.

GLADIOLUS To control Botrytis, apply Botran at the rate of 2 1/2 to 3 1/2 lbs. per acre. Do not apply to fruit that is to be eaten. Allow one day between last application and harvest.

STORAGE & DISPOSAL

Store in original container in a cool, dry place. Do not store in areas where children or animals can have access. Do not use for any purpose other than that intended. Dispose of unused material in accordance with local, state, and federal regulations.

NOTICE

Do not use for any purpose other than that intended.

WOOLFOLK CHEMICAL WORKS, INC.

MANUFACTURERS

BEST DOCUMENT AVAILABLE

FORT VALLEY, GEORGIA