

SECURITY

769-359
ACCEPTED
Oct 27 1967

EPN - 5EC

ACTIVE INGREDIENTS:

O-ethyl O-p-nitrophenyl phenylphosphonothioate 55.0%
Aromatic petroleum derivative solvent 32.1%

INERT INGREDIENTS: 12.9%

NOTE TO PHYSICIAN

Repeated therapeutic doses of Atropine may be effective. Do not use morphine. **SYMPTOMS OF POISONING** are digestive upsets, nausea, vomiting, cramps, feeling of pressure under breast bone, diarrhea, sweating, excessive salivation, restlessness, dizziness, headache, muscle tremors, twitching of the facial and eye muscles.

TREATMENT: Remove person from area of poisoning. Give two 1/100 grain atropine tablets. Do not resume operations with any organophosphorous insecticide, including EPN, until the physician has tested a sample of your blood for organophosphorous effect and tells you it is safe to do so.

and repeated as necessary for control. Borer sprays should be applied as directed by your state spray schedule, usually after harvest and at 30 day intervals for 2 to 4 sprays. Do not apply within 21 days of harvest.

PEACHES: To control Hickory Shuckworm, use 2 quarts in 500 gallons of water and apply at the rate of 250 to 500 gallons per acre. For airplane or concentrate sprayers apply at the rate of 1 to 2 quarts per acre diluted with sufficient water for coverage. Approximately 3 to 10 gallons of water per acre is the minimum amount required for coverage. Make three late applications beginning about shell hardening time which is usually about August 10. Apply a second and third spray at two week intervals. To control Pecan Weevil, use at the same rate as for Hickory Shuckworm. Applications directed against the Hickory Shuckworm will also control Pecan Weevil. Where the Pecan Weevil is the only pest to be controlled, only two applications are necessary. The first should be made when a minimum of six weevils can be larred from the branches of any one tree. This is usually about August 1 to 15. The second application should be made 10 days to two weeks later. To control Pecan Nut Casebearer, use 10 quarts in 500 gallons of water or 1 to 2 quarts per acre in sufficient water for coverage. Usually a minimum of 3 to 10 gallons of water per acre is required for concentrate sprayer. To control nut casebearer, make application at the time shuckworms have finished laying eggs. To control Pecan Nut Casebearer, use 10 quarts in 500 gallons of water or 1 to 2 quarts per acre in sufficient water for coverage. Usually a minimum of 3 to 10 gallons of water per acre is required for concentrate sprayer. To control nut casebearer, make application at the time shuckworms have finished laying eggs. To control Pecan Nut Casebearer, use 10 quarts in 500 gallons of water or 1 to 2 quarts per acre in sufficient water for coverage. Usually a minimum of 3 to 10 gallons of water per acre is required for concentrate sprayer.



POISON



DANGER: KEEP OUT OF REACH OF CHILDREN. Poisonous by skin contact, inhalation or swallowing. Rapidly absorbed through skin. Repeated exposure may, without symptoms, be increasingly hazardous. Do not get in eyes, on skin, or clothing. Do not breathe vapor or spray mist. Wear clean, heavy natural rubber gloves, rubber boots, and clean waterproof or freshly laundered protective clothing (coveralls, cap), and goggles when applying and handling. Destroy and replace gloves frequently. Wear an effective mask or respirator of a type passed by the U.S. Department of Agriculture for EPN protection. Do not contaminate feed or foodstuffs. Wash hands, arms, and face thoroughly with soap and warm water before eating or smoking. Bathe immediately after work and change all clothing. Wash clothing thoroughly with soap and hot water before re-use. In case of contact, flush skin with water. If in eyes, flush with water for 15 minutes.

PROTECT WILDLIFE: This product is highly toxic to birds, fish, and aquatic life. Do not apply to water bodies or near water. Do not apply to areas where birds, fish, or aquatic life are present. Do not apply to areas where birds, fish, or aquatic life are present.

DIRECTIONS

WARNING: This product should be used only by experienced or trained operators properly protected from the hazards involved. Use in concentrate orchard sprayers increases the hazard. Do not apply or allow to drift to areas occupied by unprotected humans or animals. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. If this product is spilled, wash down with large amounts of water. Decontaminate areas and equipment by washing with a concentrated solution of alkali and detergent and rinse with water.

DECONTAMINATE AND DESTROY CONTAINER: Rinse thoroughly with water and dump contents in a safe place. Flush and bury empty container.

PEACHES: To control Hickory Shuckworm, use 2 quarts in 500 gallons of water and apply at the rate of 250 to 500 gallons per acre. For airplane or concentrate sprayers apply at the rate of 1 to 2 quarts per acre diluted with sufficient water for coverage. Approximately 3 to 10 gallons of water per acre is the minimum amount required for coverage. Make three late applications beginning about shell hardening time which is usually about August 10. Apply a second and third spray at two week intervals. To control Pecan Weevil, use at the same rate as for Hickory Shuckworm. Applications directed against the Hickory Shuckworm will also control Pecan Weevil. Where the Pecan Weevil is the only pest to be controlled, only two applications are necessary. The first should be made when a minimum of six weevils can be larred from the branches of any one tree. This is usually about August 1 to 15. The second application should be made 10 days to two weeks later. To control Pecan Nut Casebearer, use 10 quarts in 500 gallons of water or 1 to 2 quarts per acre in sufficient water for coverage. Usually a minimum of 3 to 10 gallons of water per acre is required for concentrate sprayer. To control nut casebearer, make application at the time shuckworms have finished laying eggs. To control Pecan Nut Casebearer, use 10 quarts in 500 gallons of water or 1 to 2 quarts per acre in sufficient water for coverage. Usually a minimum of 3 to 10 gallons of water per acre is required for concentrate sprayer.

ANTIDOTE AND FIRST AID TREATMENT

INTERNAL

EXTERNAL

Wash with Soap and Warm Water.

Wash with Soap and Warm Water. Wash with Soap and Warm Water. Wash with Soap and Warm Water.

NOTICE

DISCLAIMER OF WARRANTY MERCHANTABILITY AND LIABILITY

WOOLFOLK CHEMICAL WORKS, LTD.

MANUFACTURERS

FORT VALLEY, GEORGIA