

FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 769-376
 TO ATTACHED COMMENTS

SECURITY

POWERTOX

COTTON SPRAY

EPN - METHYL PARATHION 3 - 3

ACTIVE INGREDIENTS:
 Methyl Parathion 3.0%
 Ethion 3.0%
INERT INGREDIENTS: 94.0%

U.S. EPA Reg. No. 769-376 TOTAL 100.0%

KEEP AWAY FROM HEAT AND OPEN FLAME



POISON



ANTIDOTE AND FIRST AID TREATMENT

DANGER! KEEP OUT OF REACH OF CHILDREN.

[Faint, mostly illegible text describing first aid and safety instructions]

ATROPINE IS THE EMERGENCY ANTIDOTE FOR METHYL PARATHION AND EPN POISONING.

INTERNAL

...placing a finger into the throat and giving warm salt water. ...
 ...spin of salt to a glass of water. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. **EXTERNAL:** If the material has been spilled on the skin, immediately remove patient from the vicinity of the Methyl Parathion and EPN, remove contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes. **IF WARNING SYMPTOMS APPEAR:** Administer two 1-100 grain atropine tablets immediately. NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR. See Warning Symptoms below. Keep patient calm and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, nonreactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

TREATMENT: Atropine is the specific anticholinergic agent of choice against parasympathetic nervous system activity. If there are signs of parasympathetic stimulation atropine sulfate should be administered at 15 minute intervals, in doses of 1 to 2 mg. Continue until complete normalization has occurred. Morphine is contraindicated. If chest distress persists, manage. Oxygen administration may be necessary. Advise patient to avoid sunlight for 48 hours. Repeated exposure to the water used in contact with it may cause severe conjunctivitis. If conjunctivitis occurs, irrigate eyes with sterile saline solution. Advise patient to avoid contact with the water used in contact with it for 48 hours. Advise patient to avoid contact with the water used in contact with it for 48 hours.

PROTECT WILDLIFE:

PROTECT BEES:

DISPOSAL WARNING:

DIRECTIONS

WARNING: This material is to be applied only by experienced or trained operators using airplane or power operated ground equipment. Do not apply or allow drift to reach areas occupied by unprotected humans or beneficial animals. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Keep all persons and animals out of treated area for 48 hours. If it is necessary for workers to enter treated fields within 48 hours, protective clothing should be worn. Do not apply within 5 days of harvest or of hand-picking cotton. Do not graze or feed forage from treated fields to livestock within 21 days after last application. Do not contaminate food or feed products.

COTTON: For control of overwintered Boll Weevil, Thrips, Aphids, Cotton Leafworm, Fall Armyworm, Fleahoppers, Garden Webworm, Grasshoppers, Lygus Bugs and other insects, apply at the rate of 2-3 pint per acre. For control of Boll Weevil and Bollworm in mid and late season, apply at the rate of 1-2 pints per acre. For early season Thrip control and weevil control, apply at the rate of 1-2 pints per acre. Sufficient water should be used to ensure adequate penetration. Make the first application when cotton is in the 4-5 leaf stage or equivalent. If the insecticide is used in early season, a second application should be made 7-10 days later if necessary. For control of Thrips, apply at the rate of 1-2 pints per acre. Apply at the rate of 1-2 pints per acre. Apply at the rate of 1-2 pints per acre. Apply at the rate of 1-2 pints per acre.

SOYBEANS:

NOTICE

Manufacturers

WOOLFOLK CHEMICAL WORKS, LTD. Fort Valley, Ga.