

SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY

EZE

GARDEN WEED KILLER

USE BEFORE WEEDS APPEAR
CONTAINS TREFLAN

ACTIVE INGREDIENT:
Trifluralin (a,a,a-trifluoro-2,6-dinitro-N,N-Dipropyl-p-toluidine) 1.75%
INERT INGREDIENTS: 98.25%
EPA Reg. No. 769-377 EPA Est. 769-GA-1 TOTAL 100.00%

©TREFLAN is the registered trademark for Elanco Product Company, Trifluralin.

STATEMENT OF PRACTICAL TREATMENT
In case of contact, immediately flush eyes or skin with plenty of water.
SEE ADDITIONAL PRECAUTIONARY STATEMENTS BELOW

ACCEPTED
AUG 2 1976
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under the FIFRA
EPA Reg. No. 769-377
NET CONTENTS
5 POUNDS (containing)



KEEP OUT OF REACH OF CHILDREN
CAUTION!

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS & DOMESTIC ANIMALS
CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes, or clothing.

ENVIRONMENTAL HAZARD
This pesticide is toxic to fish. Use with care when applying in areas adjacent to any body of water. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

WOOLFOLK CHEMICAL WORKS, INC.
MANUFACTURERS
FORT VALLEY, GEORGIA

DIRECTIONS FOR USE
GENERAL CLASSIFICATION

It is a violation of federal law to use this product in a manner inconsistent with its labeling.
PRECAUTIONS: This product is not recommended for use on muck soils. Do not apply to any vegetable crop not listed on the label. Do not apply to Ajuga, Myrtle, Pachysandra and other ornamental groundcovers as injury may result. Use only at the rate recommended; overtreatment may cause injury. Applications made according to directions and under normal growing conditions will not harm treated plants. Seedling disease, cold weather in combinations with deep planting and excessive moisture, high salt concentration, and drought may weaken crop seedlings and increase the possibility of damage from this product. Under these conditions, delayed crop development may result.

HOW TO APPLY: On garden soils to be planted later, thoroughly prepare the soil, break up all clods, remove all weeds and grasses and smooth the surface. Apply the recommended rate of EZE Garden Weed Killer over the surface of the area to be treated using a granular applicator or seeder. Immediately after application incorporate the herbicide into the soil to a depth of 2 to 3 inches using a roto-tiller or rake. Incorporation should be done in two directions, one at a right angle to the other, to insure more uniform distribution of the herbicide. Shallow cultivation during the season will not reduce weed control with EZE Garden Weed Killer. On established plantings, remove all existing grasses and weeds, then apply the recommended rate uniformly to the soil surface. Immediately work into the soil surface with a roto-tiller or rake being careful not to disturb the roots of the established plants.

RATES OF APPLICATION
FOR VEGETABLES

RECOMMENDED FOR USE ON Drybeans, Southern Peas (cowpeas, field peas, blackeyes peas), Snap Beans, Lima Beans, Okra and transplants of Tomatoes, Peppers, Broccoli, Brussels Sprouts, Cabbage, Cauliflower: Apply from 6 weeks before planting up to day of planting. Except on Snap Beans and Lima Beans apply at the following rates. Use 1 pound to 1500 square feet on sandy and sandy loam (light) soils. Use 1 pound to 1000 square feet on loam (medium) soils. Use 1 pound to 750 square feet on silt and clay (heavy) soils. On sandy soils containing 2-5 per cent organic matter apply 1 pound per 750 square feet. On Snap Beans and Lima Beans apply 1 pound to 1500 square feet on sandy, sandy loam or loam soils and 1 pound to 1000 square feet on silt and clay soils.

RATES OF APPLICATION
FOR ORNAMENTAL SHRUBS, TREES AND FLOWERS

RECOMMENDED FOR USE ON Established Shrubs and Trees, and these Flowers: Ageratum, Aster, Balsam, Calendula, California Poppy, Calliopsis, Carnation, Cosmos, Daffodils, Dahlia, Forget-Me-Not, Four O'Clocks, Gaillardia, Gladiolus, Hollyhock, Lobelia, Marigold, Morning Glory, Nasturtium, Periwinkle, Petunia, Portulaca, Shasta Daisy, Snapdragon, Storks, Sunflower, Sweet Alyssum, Sweet Pea, Tulips, Zinnia. Apply at the rate of 1 pound to 750 to 1000 square feet in the spring around shrubbery and trees and in flower beds after plants have become established.

WEEDS CONTROLLED
ANNUAL GRASSES

Crabgrasses	Stinkgrass
Barnyardgrass (watergrass)	Bromegrass
Foxtails (including giant foxtail)	Brachiaria
Johnsongrass (from seed)	Sandbur
Goosegrass	Jungle-Rice
Wild Cane (shattercane)	Sprangletop
Cheat	

ANNUAL BROADLEAVES

Pigweeds (spiny, redroot)	Purslane
Carelessweed	Florida Pusley
Lambsquarters	Knotweed
Carpetweed	Stinging Nettle
Russian Thistle	Goosefoot
Kochia	Chickweed

WILL NOT CONTROL certain resistant weeds such as Cocklebur, Velvetleaf, Jimsonweed, Ragweed, Venice Mallow, Nutgrass and established Bermuda and Johnsongrass.

STORAGE & DISPOSAL

STORAGE: Do not contaminate water, food, feeds, other pesticides, fertilizer or seeds.

DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

NOTICE: The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.