

SECURITY

1967 05

WOLVOLK CHEMICAL WORKS, LTD.
KILLER

NET WT. 2500

APPLY IN THE SPRING BEFORE
CRABGRASS COMES UP

CONTAINS NEW **BENEFIN** (R)

FOR LONG LASTING CRABGRASS CONTROL

K I L L S C R A B G R A S S SO YOU NEVER SEE IT

ACTIVE INGREDIENT:

N-butyl-N-ethyl-a,a,a-trifluoro-2,6-dinitro-	
B -toluidine	1.15%
INERT INGREDIENTS	98.85%
U.S.D.A. Reg. No. 769-	TOTAL 100.00%

CAUTION: Keep out of reach of children. Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. In case of contact, flush with water. Do not contaminate foodstuffs or feeds.


Do not let livestock graze treated areas. Do not feed forage from treated areas to livestock. Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes.

BENEFIN is the trademark for Elanco Products Co. benefin

WOLVOLK CHEMICAL WORKS, LTD.

Manufacturers

Fort Valley, Georgia



DIRECTIONS FOR USE

Security Crabgrass Killer is a selective herbicide for the pre-emergence control of crabgrass and other annual weeds in lawns, golf courses, and other areas where crabgrass is a problem.

- | | |
|------------------------|-----------|
| CRABGRASS (Large) | CRABGRASS |
| CRABGRASS (Small) | CRABGRASS |
| CRABGRASS (Very Small) | CRABGRASS |
| CRABGRASS | CRABGRASS |

PRECAUTIONS: Apply Security Crabgrass Killer to the turf prior to seed germination. It will not control established grass weeds. It should not be used on lawns or other areas where crabgrass is not to be applied in the spring to bluegrass, fescue, or other cool season grasses. Do not apply to newly sprigged areas of bermudagrass, St. August, or centipedegrass until these turfs are well established. Inhibition may result. Rates over 7 1/2 pounds to 2500 square feet may thin fine-leaved fescues.

GRASS WEEDS CONTROLLED: Smooth crabgrass, hairy woodgrass, yellow foxtail, (barnyard grass), goosegrass, yellow and green foxtail.

WHEN TO APPLY: Apply Security Crabgrass Killer to the turf in early spring before germination of annual grass weeds. This varies by region and climatic zone. Consult local authorities for accurate germination dates in your area. Application should be made approximately March 1 in the latitude of north Florida and the Gulf Coast area; March 15 from the coastal area of North Carolina through central South Carolina, Georgia, Alabama, Mississippi, and northern Louisiana; April 1 in central and western North Carolina, except the mountain area, and southern Tennessee, except the mountain area, and southern Missouri and northern Arkansas; April 15 in Eastern, southern Indiana and southern Illinois and northern Missouri. In regions where crabgrass and other annual weed germinates in late spring or early fall, a second application of Security Crabgrass Killer at the recommended rate may be made during the fall or the application for continued weed control.

RATE OF APPLICATION: On all turf grasses listed, except in areas where a specific rate is given, 10 pounds to 2500 square feet.

Apply in a uniform manner using a granular applicator. Do not apply to wet turf. Do not apply to areas where the turf is dormant. Do not apply to areas where the turf is being renovated. Do not apply to areas where the turf is being overseeded. Do not apply to areas where the turf is being sodded. Do not apply to areas where the turf is being repaired.

PERSONAL TRAINING: Dispose of waste pesticide and containers by
following the instructions on the label.
Do not use for anything other than the intended purpose.
Do not use if the label is missing or illegible.
Do not use if the container is damaged or leaking.
Do not use if the product is expired.