

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
DANGER

Fatal if swallowed. Do not breathe dust. Avoid contact with skin or clothing. Wash thoroughly after use. Wash equipment and clothing after application. Keep children and pets off treated areas until this material has been watered in well and grass is dry. Do not allow domestic animals to graze treated areas.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to wildlife. Use with care when applying to areas frequented by wildlife. Keep out of lakes, ponds, and streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

Read and follow all directions on label. Use this product in accordance with its labeling.

SECURITY TURF-CAL SPRAY is a fast-acting emergency crabgrass control in established turf, such as fairways, parks, golf courses, and lawns located on industrial sites. Do not use on home lawns. **SECURITY TURF-CAL SPRAY** will also control POA ANNUA (Annual Bluegrass), chickweed, and soil insect grubs (Japanese Beetle, Asiatic Beetle, and white grub). The recommended rate for control of these soil insect grubs is 3 1/2 lbs. per 1000 sq. ft. of lawn area. When a rate of either 5 or 7 lbs. per 1000 sq. ft. is used, it will control (refer to table below) control of grubs is site specific.

TIME OF APPLICATION

SPRING: Apply after ground is free from frost and well before crabgrass and Poa annua germinate. March and April are the best months for treatment. Applications made later may not be fully effective that year.
EARLY FALL: To control crabgrass and Poa annua the following year, apply well before ground freezes.

APPLICATION AND AMOUNT TO USE: Prepare turf free of weeds, mowing if necessary, so spray reaches soil surface. Measure area to be treated and uniformly spray required amount. Insufficient water for thorough coverage. Cross treating with one half required amount in each direction will give best results. If rain is not expected immediately after application, sprinkle turf thoroughly. Apply as follows:

CRABGRASS AND POA ANNUA CONTROL

Type of Grass	Type of Soil	Amount per 100 sq. ft.	When to apply
Kentucky, Meyer Bluegrass, Bermuda and Paspalum spp.	Moist	5 lbs.	Spring or Early Fall
Red Fescue and Bentgrass	Moist	7 lbs.	Spring or Early Fall

BEST DOCUMENT AVAILABLE

Manufacturers

WOOLFOLK CHEMICAL WORKS, INC.

Fort Valley, Ga.

SECURITY

SPRAY

ACTIVE INGREDIENT

1,1-Dichloro-2,2-bis(4-chlorophenyl)ethane


INERT INGREDIENTS

Water, 99.9%

Other ingredients, 0.1%

Total weight, 100%

BEST DOCUMENT AVAILABLE



KEEP OUT OF REACH OF CHILDREN.
STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Give 1 glass of water immediately and administer 15 to 30 cc of 1% solution of sodium bicarbonate. Do not induce vomiting. If unconscious, give 1 glass of water and 15 to 30 cc of 1% solution of sodium bicarbonate. **CALL A PHYSICIAN IMMEDIATELY.**
IF IN EYES OR SKIN: Flush with plenty of water. If in eyes, flush for 15 minutes. If on skin, wash with soap and water.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

REAPPLICATION: After treating in the fall and again in spring at 1000 sq. ft. apply annual rate of 2 1/2 to 3 1/2 pounds per 1000 sq. ft. to maintain control of crabgrass and annual bluegrass.

WHERE POA ANNUA IS A MAJOR PROBLEM: SPRING TREATMENT Apply 4 to 5 lbs. per 1000 sq. ft. Repeat treatment again before April 15 and October 15. To maintain soil toxicity, make another application at 1000 sq. ft. of Turf-Cal Spray.

PRECAUTIONS: Water Turf-Cal Spray in well right after application. Do not get material in contact with soil and prevent any temporary watering. Do not overlap areas when applying nor skip any areas. Keep away from lawn to that if spilled, lawn will not be harmed. Do not exceed rate of application recommended. Inadequate watering may cause some temporary scorch. Do not apply to Dichondra or St. Augustine lawns. Avoid contact with flowers and shrubs and application in the shaded or hot zones. Do not apply Turf-Cal Spray to heavily shaded areas until grass has become well established and has received a second cutting. Do not use a high phosphate fertilizer while program is in force. To help desirable grasses thrive, a high nitrogen fertilizer may be applied in spring and fall as needed. Apply fertilizer in spring and fall, but not at the same time as Turf-Cal Spray. Leave an interval of at least several months between applications of Turf-Cal Spray.

SPECIAL NOTES: Kentucky, Meyer Bluegrass, Zoysias, Bermuda and Paspalum are more tolerant of arsenic than Fescues and Bentgrass.

TURF RENOVATION: Vertical cutting and overseeding should be employed before or even during building of toxicity. Good vegetative improvement may also be desired.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not use or store in or around the home. Do not contaminate water, food or feed by storage or disposal. Open burning prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL: Do not pour mixture or residue that can be collected. Material which cannot be collected should be disposed of according to local, state and federal laws and by local, state or federal disposal authority.

CONTAINER DISPOSAL: Do not reuse. State or local disposal authority should be contacted.

NOTICE: This product is sold with no warranty of merchantability nor with any warranty which extends beyond the description on this label. 977190