

**ACCEPTED**  
**JUN 28 1985**  
 Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act,  
 as amended, for the pesticide  
 registered under  
 EPA Reg. No. 769-413

769-413 PM 12  
 131



# Bait-A-Tox S

**ALFALFA:** Apply 30 pounds per acre. Must be applied at least 3 days prior to harvest.  
**COTTON:** Apply 30 pounds per acre. Must be applied at least 7 days prior to harvest.

### STORAGE AND DISPOSAL

**STORAGE:** Store pesticide in original container and place in a locked storage area. Opened pesticide containers should be securely resealed prior to storage. Protect from penetration of moisture, excessive heat or cold. Avoid cross-contamination with other pesticides or fertilizers. Use caution when storing pesticides in stacks.

**PESTICIDE DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### NOTICE

#### DISCLAIMER OF WARRANTY MERCHANTABILITY AND LIABILITY

If for any reason this product is not used, applied, stored, transported or disposed of in accordance with the instructions set forth on this label, then and thereafter Security Chemical Company terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness.

NET WEIGHT: 50 POUNDS

**SECURITY CHEMICAL COMPANY**  
 MANUFACTURERS ATLANTA, GEORGIA

**ACTIVE INGREDIENT:**  
 Carbaryl (1-naphthyl N-methylcarbamate) . . . . . 5.0%

**INERT INGREDIENTS:** . . . . . 95.0%

**TOTAL** . . . . . 100.0%

EPA Reg. No. 769-413                      EPA Est. 769-GA-1

**KEEP OUT OF REACH OF CHILDREN  
 AND ANIMALS!**

### CAUTION!

(See additional precautionary statements on the side panel.)

#### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention. **IF INHALED:** Remove victim to fresh air. **IF ON SKIN:** Wash with soap and water. Remove any contaminated clothing. **IF IN EYES:** Flush eyes with water.

**NOTE TO PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM opiates or cholinesterase inhibiting drugs.

#### PRECAUTIONARY STATEMENTS

**CAUTION:** May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

#### ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water

or wetlands. Do not contaminate water by cleaning of equipment or disposal of waste.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels. Applications may be made with either ground equipment or aircraft. Repeat as necessary. Bait-A-Tox will control cutworms, armyworms, crickets, darkling ground beetles, grasshoppers and sowbugs.

#### VEGETABLES

**CUCUMBERS, SQUASH, MELONS:** Apply 20 pounds per acre. May be applied on day of harvest.

**PEASE:** Apply 30 pounds per acre. May be applied on day of harvest. (Cowpeas: Must be applied at least 3 days before harvest.)

**BEANS, CARROTS, EGGPLANTS, OKRA, PEPPERS, POTATOES, and SWEET CORN:** Apply 40 pounds per acre. May be applied on day of harvest.

**ASPARAGUS, STRAWBERRIES:** Apply 40 pounds per acre. Do not apply within one day of harvest. Do not repeat application on asparagus within 3 days.

**BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, HEAD LETTUCE, GARDEN BEETS (Roots), HORSE RADISH, PARSNIPS,**

**RADISHES, RUTABAGAS and TURNIPS (Roots):** Apply 40 pounds per acre. Do not apply within 3 days of harvest.

**SUGAR BEETS, COLLARDS, ENDIVE (Escarole), GARDEN BEETS (Tops), KALE, LEAF LETTUCE, PARSLEY, SPINACH, SWISS CHARD and TURNIPS (Tops):** Apply 40 pounds per acre. Do not apply within 14 days of harvest.

**PEANUTS:** Apply 20 pounds per acre. May be applied on day of harvest.

**CORN:** Apply 40 pounds per acre. May be applied on day of harvest.