

(1 fl. oz.) in 3 gallons of water. Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to penetrate dense foliage.

**FRUITS:** Apples - to control Codling Moths, Mites, Aphids, Leaf Rollers, and Scale Crawlers. Peaches - to control Mites, Aphids, Leaf Rollers, and Scale Crawlers. Oranges - to control Mites, Aphids, and Scale Crawlers. Plums - to control Mites, Aphids, Leaf Rollers and Scale Crawlers. Grapes - to control Mites, Aphids, Leaf Rollers and Scale Crawlers. Cherries - to control Mites, Aphids and Scale Crawlers. Strawberries - to control Mites, Aphids and Leaf Rollers. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water. Apply as a full cover spray. Start spraying after blooming and repeat at 10 to 14 day intervals if needed. Do not spray fruit within 21 days of picking.

**VEGETABLES:** Tomatoes - to control Aphids, Leaf Miners. Carrots - to control Aphids. Potatoes, Beans - to control Aphids, Leaf Miners, Bean Beetles. Cucumbers, Cantaloupe, Peppers - to control Aphids, Leaf Miners, Cucumbers.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

# IMAZINON 2EC

FOR USE ON LAWNS, FLOWERS, ORNAMENTALS, VEGETABLES, FRUITS, OUTDOOR AREAS

## ACTIVE INGREDIENTS:

O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate .... 25.0%  
Aromatic Petroleum Derivative Solvent ..... 55.7%

## INERT INGREDIENTS:

..... 19.3%

E.P.A. Reg. No. 799-111-1

(Do not use on crops. Protective information may be obtained from the manufacturer.)

KEEP OUT OF REACH OF CHILDREN

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS

DO NOT USE ON CROPS



WOLFPOLE CHEMICAL COMPANY, INC.

Manufacturers

**WARNING!** May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not permit children or pets to go onto the sprayed grass until the spray has completely dried. Keep out of reach of domestic animals. Do not store or use near heat or open flame. Do not contaminate ornamental fishponds.

**PROTECT WILDLIFE:** This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**PROTECT BEES:** This product is highly toxic to bees exposed to direct contact with treated areas. Protective information may be obtained from the manufacturer.

**INSTRUCTIONS:**

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

**FRUITS:** Apples - to control Aphids, Leaf Miners, Red Worms, Scale Crawlers, Pear Borer, Pear Rust, Red Bugs, Soap Scale, etc. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. of surface. Do not spray within 21 days of picking. Do not spray within 14 days of harvesting potatoes, or 14 days of harvesting other crops.

ACCEPTED

JUL 3 1975

U.S. Dept. of Agriculture  
Federal Insecticide,  
Fungicide, and Rodenticide Act  
Registration of this pesticide  
is complete for use as a  
In A. Reg. No.

1. Mix 2 tbsps. of water. Apply as a foliar spray. Thoroughly wet the undersides of the leaves. Repeat at 7-10 day intervals.

**FRUITS:** Apples - to control Chewing Moths, Mites, Aphids, Leaf Rollers, and Scale Crawlers. Peaches - to control Mites, Aphids, Leaf Rollers, and Scale Crawlers. Oranges - to control Mites, Aphids, and Scale Crawlers. Plums - to control Mites, Aphids, Leaf Rollers, and Scale Crawlers. Grapes - to control Mites, Aphids, Leaf Rollers, and Scale Crawlers. Cherries - to control Mites, Aphids, and Scale Crawlers. Strawberries - to control Mites, Aphids, and Leaf Rollers. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water. Apply as a foliar spray. Start spraying after blooming and repeat at 7-10 day intervals if needed. Do not spray fruit within 21 days of picking.

**VEGETABLES:** Tomatoes - to control Aphids, Leaf Miners. Carrots - to control Aphids. Potatoes, Beans - to control Aphids, Leaf Miners, Bean Beetles. Cucumbers, Cantaloupe, Peppers - to control Aphids, Leaf Miners, Cucumber Beetles. Onions - to control Aphids. Cabbage - to control Aphids, Leaf Miners, Cabbageworms. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 1,000 sq. ft. Generally application at 7-10 day intervals may be required. Do not apply within one day of picking tomatoes, 35 days of harvesting potatoes, or 14 days of picking all other crops listed in this section.

Cabbage, Carrots, Corn, Lettuce, Peas, Pole Beans, Radishes, Red Beets, Snap Beans, Tomatoes, Turnips - to control cutworms (Surface and Subterranean), Wireworms, Root Maggots, Garden Symphylans, Mole Crickets. Apply 15 fl. oz. per 1,000 sq. ft. as a soil treatment in sufficient water to obtain thorough coverage. Make application to soil surface as a broadcast spray prior to planting. Thoroughly mix the chemical into the top 4-8 inches of soil.

**NOTE:** Apply not more than 2 tbsps. (1 fl. oz.) to 1350 sq. ft. of soil surface where beans, carrots, cucumbers, onions and cabbage will be planted and 2 tbsps. to 2700 sq. ft. of soil surface where peppers will be planted.

**OUTSIDE NUISANCE PESTS:** Flies, Ants, Cockroaches, Boxelder Bugs, Clover Mites, Spiders, Ticks, Chiggers, Earwigs. Apply 8 fl. oz. per 3 gallons of water. Thoroughly spray window frames, doors, sills, screens, porch and patio walls and ceiling, foundations, walks, garbage cans, and into cracks and other places where insects may hide. Do not treat plants with this dosage rate.

**DISPOSAL WARNING:** Dispose of waste pesticide and spillage by burying in non-crop land away from water supplies. Do not re-use containers; destroy when empty.

#### NOTICE

The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label. 370314

#### FOR USE ON LAWNS, FLOWERS, ORNAMENTALS, VEGETABLES, FRUITS, OUTDOOR AREAS

##### ACTIVE INGREDIENTS:

D-D-ethyl (O-2-isopropyl-6-methyl-4-pyridinyl) phosphorothioate 25.0%  
Aromatic Petroleum Derivative Solvent 55.0%

**INERT INGREDIENTS:** 19.3%

E.P.A. Reg. No. 769-317 E.P.A. Est. No. 769-GA-1 TOTAL 100.0%

(R) Diazinon, a registered trademark of CIBA-GEIGY Corporation.

**CAUTION: COMBUSTIBLE MIXTURE. DO NOT USE IN THE HOME.**

**WARNING!**  
KEEP OUT OF REACH  
OF CHILDREN.

May be fatal  
if swallowed.  
See side panel  
for additional  
precautions.



NET CONTENTS  
ONE GALLON

**WOOLFOLK CHEMICAL WORKS, INC.**  
Manufacturers Ft. Valley, Ga.

#### WARNING!

May be fatal if swallowed. May irritate the eyes, nose, throat, and skin. Do not breathe spray mist. Avoid contact with skin. Wash thoroughly after handling. Avoid contact with food and foodstuffs.

Use on household pets or humans. Do not permit children or pets to enter the sprayed grass until the spray has completely dried. Keep out of reach of nesting animals. Do not store or use near heat or open flame. Do not use near ornamental fishponds.

#### PROTECT WILDLIFE:

This product is highly toxic to birds. Do not feed or treat areas may be used. Keep out of lakes, streams, ponds, or not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**PROTECT BEES:** This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

#### DIRECTIONS

**LAWNS:** Bluegrass, Merion, Fescue, St. Augustine, Dichondra, Bent, Bermuda. To control Chinch Bugs\*, Sod Webworms, Bermuda Mites, Ants, Ticks, Earwigs, Lawn Billbugs. Apply 2 tbsps. (1 fl. oz.) in 3 gallons of water per 125 sq. ft. or 8 fl. oz. in 24 gallons of water per 1,000 sq. ft. To control Millipedes - Apply 4 tbsps. (2 fl. oz.) in 3 gallons of water per 125 sq. ft. or 16 oz. in 24 gallons of water per 1,000 sq. ft.

Thoroughly wet down grass a few hours before applying. For Lawn Billbugs, water lawn thoroughly after treatment. \*In the north, if Chinch Bugs are the only insect problem, a lower rate of application can be used; 2 tbsps. in 3 gallons of water per 250 sq. ft.

To control Rhodes - Grass Scale Crawlers. Apply 2 tbsps. (2 fl. oz.) in 3 gallons of water per 100 sq. ft. or 10 oz. in 30 gallons of water per 1,000 sq. ft. Apply when Scale Crawlers first emerge.

**ROSES AND FLOWERS:** Roses, Azaleas, Chrysanthemums, Gladioli, Carnations. To control Aphids, Mites, Leafhoppers, Thrips - Apply 2 tbsps. (1 fl. oz.) per 3 gallons of water. Apply as a thorough foliar spray. Spray underside of leaves. Wet plants to the dripping point.

**TREES AND SHRUBS:** Oak - To control Aphids and Scale Crawlers. Maple - To control Aphids. Locust - To control Aphids. Holly and Boxwood - To control Mites, Scale Crawlers, and Leaf Miners. Pine - To control Aphids and Scale Crawlers. Spruce - To control Aphids, Scale Crawlers, and Bagworms. Willow - To control Aphids, Scale Crawlers, and Leaf Miners. Apply 2 tbsps.