



Diazinon® 2-EC

FOR USE ON LAWNS, FLOWERS, ORNAMENTALS,
VEGETABLES, FRUITS, OUTDOOR AREAS.

ACTIVE INGREDIENTS:

O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)
phosphorothioate 25.0%

Aromatic Petroleum Derivative Solvent 55.7%

INERT INGREDIENTS: 19.3%

Total 100.0%


E.P.A. Reg. No. 769-417 E.P.A. Est. No. 769-GA-1

®Diazinon is a registered trademark of CIBA-GEIGY CORPORATION.

**THIS PRODUCT CONTAINS 2 LBS. OF
DIAZINON® PER GALLON.**

**KEEP OUT OF REACH OF CHILDREN.
WARNING**

(See additional precautionary statements on the back panel.)



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS WARNING

May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not permit children or pets to go onto sprayed areas until the spray has completely dried. Keep out of reach of domestic animals. Do not contaminate ornamental fishponds. DO NOT USE IN THE HOME.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PROTECT BEES: This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

LAWNS: Bluegrass, Merion, Fescue, St. Augustine, Dichondra, Bent, Bermuda. To control Chinch Bugs*, Sod Webworms, Bermuda Mites, Ants, Ticks, Earwigs. Lawn Billbugs, apply 2 tbsps. (1 fl. oz.) in 3 gals. of water per 125 sq. ft. or 8 fl. oz. in 24 gals. of water per 1,000 sq. ft. To control Millipedes, apply 4 tbsps. (2 fl. oz.) in 3 gals. of water per 125 sq. ft. or 16 oz. in 24 gals. of water per 1,000 sq. ft. Thoroughly wet down grass a few hours before applying. For Lawn Billbugs, water lawn thoroughly after treatment. In the North, if Chinch Bugs are the only insect problem, a lower rate of application can be used: 2 tbsps. in 3 gals. of water per 250 sq. ft. To control Rhodes Grass Scale Crawlers, apply 2 tbsps. (1 fl. oz.) in 3 gals. of water per 100 sq. ft. or 10 oz. in 30 gals. of water per 1,000 sq. ft. Apply when Scale Crawlers first emerge.

FIRE ANTS: To aid in the control of fire ants in lawns and other recreational areas, apply 1 fl. oz. (2 tbsps.) in a minimum of 1 gal. of water as a drench. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Larger quantities of solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted product as a gentle rain at the rate of 1 gal. per mound. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness. For best results, apply in cool weather, 65-80°F, or in early morning or late evening hours. Treat new mounds as they appear.

ROSES AND FLOWERS: Rose, Azaleas, Chrysanthemums, Gladiolus, Carnations. To control Aphids, Mites, Leafhoppers, Thrips, apply 2 tbsps. (1 fl. oz.) per 3 gals. of water. Apply as a thorough foliar spray. Spray underside of leaves. Wet plants to the dripping point.

TREES AND SHRUBS: Oak - To control Aphids and Scale Crawlers. Maple - To control Aphids, Locust - To control Aphids. Holly and Boxwood - To control Mites, Scale Crawlers, and Leaf Miners. Pine - To control Aphids and Scale Crawlers. Spruce - To control Aphids, Scale Crawlers, and Bagworms. Willow - To control Aphids, Scale Crawlers, and Leaf Miners. Apply 2 tbsps. (1 fl. oz.) in 3 gals. of water. Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to penetrate dense foliage.

FRUITS: Apples - To control Codling Moths, Mites, Aphids, Leaf Rollers, and Scale Crawlers. Peach - To control Mites, Aphids, Leaf Rollers, and Scale Crawlers. Oranges - To control Mites, Aphids, and Scale Crawlers. Plums - To control Mites, Aphids, Leaf Rollers and Scale Crawlers. Grapes - To control Mites, Aphids, Leaf Rollers and Scale Crawlers. Cherries - To control Mites, Aphids, and Scale Crawlers. Strawberries - To control Mites, Aphids, and Leaf Rollers. Apply 2 tbsps. (1 fl. oz.) in 3 gals. of water. Apply as a full cover spray. Start spraying after blooming and repeat at 10 to 14 day intervals if needed. Do not spray fruit within 21 days of picking.

VEGETABLES: Tomatoes - To control Aphids, Leaf Miners. Carrots - To control Aphids, Potatoes, Beans - To control Aphids, Leaf Miners, Bean Beetles, Cucumbers, Cantaloupe, Peppers - To control Aphids, Leaf Miners, Cucumber Beetles, Onions - To control Aphids, Cabbages - To control Aphids, Leaf Miners, Cabbageworms. Apply 2 tbsps. (1 fl. oz.) in 3 gals. of water per 1,000 sq. ft. Generally application at 7-10 day intervals may be required. Do not apply within one day of picking tomatoes, 35 days of harvesting potatoes, or 14 days of picking all other crops listed in this section.

Cabbage, Corn, Lettuce, Peas, Pole Beans, Radishes, Red Beans, Snap Beans, Tomatoes, Turnips - To control Cutworms (Surface and Subterranean), Wireworms, Root Maggots, Garden Symphylans, Mole Crickets. Apply 15 fl. oz. per 1,000 sq. ft. as a soil treatment in sufficient water to obtain thorough coverage. Thoroughly mix the chemical into the top 4-8 inches of soil.

NOTE: Apply not more than 2 tbsps. (1 fl. oz.) to 150 sq. ft. of soil surface where beans, carrots, cucumbers, onions and cabbage will be planted and 2 tbsps. to 2700 sq. ft. of soil surface where peppers will be planted.

OUTSIDE NUISANCE PESTS: Flies, Ants, Cockroaches, Boxelder Bugs, Clover Mites, Spiders, Ticks, Chiggers, Earwigs. Apply 8 fl. oz. per 3 gals. of water. Thoroughly spray window frames, doors, eaves, screens, porch and patio walls and ceiling, foundations, walks, garbage cans, and into cracks and other places where insects may hide. Do not treat plants with this dosage rate.

DISPOSAL

Do not reuse container. Wrap container and put in trash collection.

NOTICE: Seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

ONE FL. QUART

Manufactured by
WOOLFOLK CHEMICAL WORKS, INC. Fort Valley, Ga.

Back panel
over label

769-417

07/26/1979

1/1