


WOOLFOLK CHEMICAL WORKS, LTD

ROSE and FUNGICIDE

Contains Benomyl

ACTIVE INGREDIENTS:
Benomyl
[Methyl 1-(butylcarbamoyl)
2-benzimidazole-
carbamate] 50%
INERT
INGREDIENTS 50%
U.S. Pat. Per d. n.
EPA Reg. No. 769-419

NET WEIGHT
2 OUNCES



Controls Certain
Diseases of
Roses, Flowers,
Ornamentals,
and Turf.

CAUTION:
Keep out of reach
of children
(See back label for
additional precautions)

270127

FORT VALLEY, GEORGIA 31030

CAUTION: Keep out of reach of children. May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, nose, and clothing. Wash thoroughly after using. Keep away from fire and sparks. In case of contact flush skin or eyes with plenty of water. For eyes get medical attention.

DIRECTIONS: Security Rose and Turf Fungicide has both curative and protective action and is highly effective for the control of many fungus diseases. It is a wettable powder and must be mixed with water and applied as a spray. Addition of a surfactant to the spray mixture enhances curative action of the fungicide and improves the distribution of the spray on hard to wet plants such as roses. It may be applied with compressed air sprayers, power sprayers and certain hose sprayers.

ROSES: To control Powdery Mildew and Black Spot use one tablespoonful to 2 gals. water. Begin application when disease first appears and repeat at 10- to 14-day intervals throughout the growing season; shorten intervals during humid, rainy weather.

FLOWERS, ORNAMENTALS: To control Powdery Mildew and Botrytis Grey Mold use one tablespoonful to 2 gals. water. Begin application when disease first appears and repeat at 10- to 14-day intervals as long as needed.

IMPORTANT: Store in a cool, dry place. Do not use if container is damaged. Do not use if contents are clumpy. Do not use if contents are discolored. Do not use if contents are otherwise changed. Do not use if contents are otherwise changed. Do not use if contents are otherwise changed. Do not use if contents are otherwise changed.

LAWN AND TURF GRASSES: To control Dollar Spot and Large Brown Patch, apply at the rate of 2 oz. (12 tablespoonfuls) per 1000 sq. ft. in sufficient water to obtain thorough coverage, usually 2 to 5 gals. per 1000 sq. ft. Apply at earliest appearance of disease and continue at 10- to 14-day intervals as long as needed. When conditions are unusually favorable for development of disease, reduce interval to 5 to 7 days. Do not graze or feed clippings from treated areas to livestock.

DISPOSAL WARNING: Dispose of waste pesticide and container in a safe place. Never re-use container.

NOTICE: The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

Sold By
WOOLFOLK CHEMICAL WORKS, LTD.
Fort Valley, Georgia

