

FEDERAL PESTICIDE CONTROL ACT
EPA REGISTRATION NUMBER
EPA REG. NO. 767-423

SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY

DO NOT ALLOW SPRAY TO COME IN CONTACT WITH FOLIAGE, GREEN BARK, GRAFTED UNIONS OR SCUFFED, DAMAGED OR BROKEN BARK WHEN SPRAYING AROUND TREES AND ORNAMENTALS.

DO NOT CONTAMINATE WATERS USED FOR DOMESTIC CONSUMPTION, OR BY ANIMALS, WILDLIFE AND AQUATIC LIFE, OR FOR IRRIGATION PURPOSES. DO NOT GRAZE TREATED AREAS TO LIVESTOCK.

LAWN RENOVATION: Mow lawn to about 1 inch high before treatment. Apply with either tank type or hose-end sprayer. Follow directions in the table below. Spray foliage thoroughly, preferably on a warm sunny day. Foliage will usually turn brown in about five days. Bermudagrass and other deep-rooted perennials, or partially sprayed foliage, may require a second treatment. Reapply five days later if green areas remain.

SONITE HERBICIDE

**SODIUM CACODYLATE LIQUID PLUS
SURFACTANT FOR
GENERAL POST-EMERGENT WEED CONTROL**

ACTIVE INGREDIENTS:

Sodium Cacodylate	6.74%
Dimethylarsinic Acid (Cacodylic Acid)	1.17%
INERT INGREDIENTS:	92.09%

CAUTION! KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near food or food products. Keep children and domestic animals off treated areas until this material has been washed into the soil.

ANTIDOTE: If taken internally, induce vomiting and call physician at once.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT.
GENERAL INFORMATION: Sonite Herbicide is useful for general post-emergent weed control. It is non-selective in its action. Its phytotoxic properties are quickly inactivated on contact with soil. It contains a surfactant. It is unnecessary to add any other surfactant to the spray solution. Best results are obtained on young actively growing weeds. It produces top-kill only



769-423

SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY

DO NOT ALLOW SPRAY TO COME IN CONTACT WITH FOLIAGE, GREEN BARK, GRAFTED UNIONS OR SCUFFED, DAMAGED OR BROKEN BARK WHEN SPRAYING AROUND TREES AND ORNAMENTALS.

DO NOT CONTAMINATE WATERS USED FOR DOMESTIC CONSUMPTION, OR BY ANIMALS, WILDLIFE AND AQUATIC LIFE, OR FOR IRRIGATION PURPOSES. DO NOT GRAZE TREATED AREAS TO LIVESTOCK.

LAWN RENOVATION: Mow lawn to about 1 inch high before treatment. Apply with either tank type or backpack sprayer. Follow directions in the table below. Spray foliage thoroughly, preferably on a warm sunny day. Foliage will usually turn brown in about five days. Bermudagrass and other deep-rooted perennials, or partially sprayed foliage, may require a second treatment. Reapply five days later if green areas remain.

SONITE HERBICIDE

**SODIUM CACODYLATE LIQUID PLUS
SURFACTANT FOR
GENERAL POST-EMERGENT WEED CONTROL**

ACTIVE INGREDIENTS:

Sodium Cacodylate	5.74
Dimethylarsinic Acid (Cacodylic Acid)	1.17

INERT INGREDIENTS: 92.09

CAUTION! KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near food or food products. Keep children and domestic animals off treated areas until this material has been washed into the soil.

ANTIDOTE: If taken internally, induce vomiting and call physician at once.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT.

GENERAL INFORMATION: Sonite Herbicide is useful for general post-emergent weed control. It is non-selective in its action. Its phytotoxic properties are quickly inactivated on contact with soil. It contains a surfactant. It is unnecessary to add any other surfactant to the spray solution. Best results are obtained on young actively growing weeds. It produces top-kill only

