

May 18, 1973

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
FD UNDER NO. 769-424

SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY

ditchbanks, rights-of-way, fence rows, storage yards and similar non-crop areas. Mix at a rate of 2 to 5 quarts in 100 gallons of water (4 to 10 tablespoons to 3 gallons). Spray unwanted vegetation thoroughly to just short of run-off. Any spray equipment that gives good coverage may be used. If regrowth occurs, reapply as required.

**LAWN AND ORNAMENTAL TURF:** This product is useful for control of dallisgrass, sandbur, bahiagrass, nutsedge, crabgrass, chickweed, and wood sorrel with little or no injury to good lawn grasses. On new lawns, do not treat until after three mowings. Good grasses may be temporarily discolored. Zoysia, bluegrasses and bermuda are quite tolerant. Do not use on St. Augustine, Centipede or Carpetgrass.

# CRABGRASS KILLER

**SPRAY FOR GRASSY TYPE WEEDS IN  
LAWNS & NON-CROP AREAS**

<b>ACTIVE INGREDIENT:</b>	
Monosodium Acid Methanearsonate* .....	35.6%
<b>INERT INGREDIENT:</b> .....	
64.4%	
E. P. A. Reg. No. 769-424	TOTAL 100.0%
Total Arsenic (as elemental) all in water soluble form 15.5%	
*Product contains 1 lb. MSMA per quart.	

**CAUTION!** KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near food or food products. Keep children and domestic animals off treated areas until the material has been washed into the soil.  
**ANTIDOTE:** If taken internally, induce vomiting and call physician at once.

**READ ENTIRE LABEL BEFORE USING THIS PRODUCT.**

**GENERAL INFORMATION:** This product is useful for selective post-emergent weed control, particularly for



**SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY SECURITY**

ditchbanks, rights-of-way, fence rows, storage yards and similar non-crop areas. Mix at a rate of 2 to 5 quarts in 100 gallons of water (4 to 10 tablespoons to 3 gallons). Spray unwanted vegetation thoroughly to just short of run-off. Any spray equipment that gives good coverage may be used. If regrowth occurs, reapply as required.

**LAWN AND ORNAMENTAL TURF:** This product is useful for control of dallisgrass, sandbur, bahiagrass, nutsedge, crabgrass, chickweed, and wood sorrel with little or no injury to good lawn grasses. On new lawns, do not treat until after three mowings. Good grasses may be temporarily discolored. Zoysia, bluegrasses and bermuda are quite tolerant. Do not use on St. Augustine, Centipede or Carpetgrass.

**CRABGRASS KILLER**

**SPRAY FOR GRASSY TYPE WEEDS IN LAWNS & NON-CROP AREAS**

**ACTIVE INGREDIENT:**

Monosodium Acid Methanearsonate\* ..... 35.6%

**INERT INGREDIENT:** ..... 64.4%

E. P. A. Reg. No. 769-424 TOTAL 100.0%

Total Arsenic (as elemental) all in water soluble form 16.5%

\*Product contains 1 lb. MSMA per quart.

**CAUTION!** KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near feed or food products. Keep children and domestic animals off treated areas until the material has been washed into the soil.

**ANTIDOTE:** If taken internally, induce vomiting and call physician at once.

**READ ENTIRE LABEL BEFORE USING THIS PRODUCT.**

**GENERAL INFORMATION:** This product is useful for selective post-emergent weed control, particularly for

