

NET WEIGHT

SECURITY

FOUR POUNDS

TOMATO & VEGETABLE DUST

CONTAINS SEVIN (R) AND MANEB FOR INSECT AND DISEASE CONTROL

ACTIVE INGREDIENTS:

Maleic Manganoethylenedithiocarbamate 100% 100.00
 Carbaryl (1-naphthyl N-methylcarbamate) 90% 50.00

INERT INGREDIENTS:

E. P. A. Reg. No. 7-9-437 100.00 100.00

(R) Trademark of Union Carbide Corp. for 1-naphthyl N-methylcarbamate

CAUTION! KEEP OUT OF REACH OF CHILDREN.

Harmful if inhaled, swallowed, or absorbed through the skin. Do not breathe dust. Do not take internally. Avoid skin contact. Avoid getting in eyes. Wash hands before eating. Wash all contaminated clothing with soap and water before reuse. Avoid contamination of food stuffs, feed, drugs, and water receptacles. Wash after handling.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

PROTECT WILDLIFE: This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of waste. Apply only as specified on the label.

PROTECT BEES: This product is highly toxic to bees exposed to direct treatment or residues on blossoms. Protective information may be obtained from your Cooperative Agricultural Extension Service.

STORAGE: Store in cool, dry place. Keep away from fire and sparks. Do not allow to become wet or overheated during storage. This may cause chemical changes which will impair the fungicidal effectiveness of this product.

DIRECTIONS

BEANS: To control anthracnose, downy mildew, Mexican bean beetle, bean leaf beetle, cucumber beetles, leafhoppers, flea beetles, Japanese beetle, and corn earworm. Apply at the rate of 20 pounds per acre or 1 ounce per 50 feet of row. Where Mexican bean beetle is the only problem, a rate of 10 pounds per acre is sufficient. To control squash bug, apply at 40 pounds per acre. To control Japanese beetle, apply at 40 pounds per acre or 2 ounces per 50 feet of row. For disease control, begin application when disease first appears and repeat at weekly intervals. For insect control, begin application when insects first appear and re-treat as necessary. Do not apply within 4 days of harvest.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI: To control alternaria leaf spot, downy mildew, flea beetle, harlequin bug, corn earworm, armyworm, imported cabbage worm, and diamond back moth larvae. Apply at 40 pounds per acre, or 1 to 2 ounces to 50 feet of row. Do not control cabbage loopers. For disease control, begin application as soon as disease threatens and repeat at 7 to 10 day intervals in the field. Use at 3 day intervals in plant beds. For insect control, begin application when insects first appear and re-treat as necessary. Do not apply to cabbage within 7 days of harvest or to the other crops within 7 days of harvest. Remove residue from broccoli by washing or trimming.

COLLARDS, KALE, LETTUCE, MUSTARD GREENS, SPINACH, TURNIPS (TOPS): To control alternaria leaf spot, downy mildew, white rust, flea beetle, harlequin bug, leafhopper, armyworm, imported cabbage worm, diamond-back moth larvae, corn earworm, tarnished plant bug, and stink bug. Use 20 to 40 pounds per acre or 1 to 2 ounces to 50 feet of row. For best control of corn earworm after lettuce heads begin to form, use 40 pounds per acre on a 5 to 7 day schedule. Will not control cabbage looper. For disease control, begin application as soon as disease threatens and repeat at 7 to 10 day intervals. For insect control, begin application when insects first appear and re-treat as necessary. Do not apply within 14 days of harvest. Do not apply to head lettuce within 7 days of

harvest. Apply within 14 days of harvest. Remove excess residues by trimming and washing or other effective means.

CUCUMBERS, SQUASH: To control anthracnose, downy mildew, lybium fruit rot, alternaria (Macrosporium) leaf spot, angular leaf spot, pickleworm, cucumber beetle, squash bug, and flea beetle. Apply at the rate of 20 pounds per acre or 1 ounce to 50 feet of row. For disease control, begin application when tumors begin to form at the first sign of disease. Repeat at weekly intervals. Direct dust to underside of leaves. For insect control, begin application when insects first appear and re-treat as necessary. Do not apply within 5 days of harvest.

CORN (SWEET): To control helminthosporium blight, corn earworm, European corn borer, fall armyworm, sap beetle, and Japanese beetle. Apply at the rate of 30 to 40 pounds per acre or 2 ounces to 50 feet of row. For European corn borer, consult your state Agricultural Experiment Station for proper timing of treatments. For budworm damage in the whorl, apply to entire plant and repeat as necessary. For insects attacking silks and ears, apply at 3 to 5 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. For disease control, begin application when plants are 6 inches high and repeat at 5 to 7 day intervals until harvest. For insect control, begin application when insects first appear and re-treat as necessary. Do not feed treated forage to livestock.

CARROTS: To control alternaria and necrotic leaf spot, flea beetles, leafhoppers, spittlebugs, lygus and stink bugs. Apply at the rate of 20 to 40 pounds per acre or 1 to 2 ounces to 50 feet of row. For disease control, begin application when plants are 6 weeks old and repeat at 7 to 10 day intervals. For insect control, begin application when insects first appear and re-treat as necessary.

TOMATO, EGGPLANT, PEPPER: To control anthracnose, late and early blight, alternaria, phytophthora blight or fruit rot, necrotic leaf spot, gray leaf spot, ripe rot, cladosporium leaf mold, gray leaf spot (Stenphythium), septoria leaf spot, tomato fruitworm, fall armyworm, flea beetles, leafhoppers, tomato hornworm, European corn borer and stink bug. Apply at the rate of 20 to 40 pounds per acre. For disease control, begin application at first cluster and repeat at 7 to 10 day intervals or oftener during severe late blight conditions. To avoid damage, do not use on tender young greenhouse tomato plants. For insect control, begin application when insects first appear and re-treat as necessary. Do not apply within 5 days of harvest.

POTATO: To control early and late blights, Colorado potato beetle, leafhoppers, and flea beetles. Apply at 20 to 40 pounds per acre or 1 to 2 ounces to 50 feet of row. Where Colorado potato beetle is the only problem, a rate of 10 pounds per acre is sufficient. To control lygus bug, apply at 40 pounds per acre. For disease control, begin application when plants are 2 to 6 inches high. Repeat at 5 to 10 day intervals, more often under severe late blight conditions. For insect control, begin application when insects first appear and re-treat as necessary.

DISPOSAL WARNING

Dispose of waste by burying in noncrop lands away from water supplies. Containers should be disposed of by burying with wastes, not by burning. Stay out of smoke. Never re-use container.

NOTICE

The seller makes no express warranty, merchantability nor any other warranties which extend beyond the description on this label.

WOOLFOLK CHEMICAL WORKS, INC.

Manufacturers

Fort Valley, Ga.

ACCEPTED

APR 24 1973

U.S. DEPT. OF AGRICULTURE
 ARIZONA STATE UNIVERSITY
 TUCSON, ARIZONA
 TO THE DIRECTOR