

SECURITY

TOX - SOL - 8

CONTAINS 8 POUNDS TOXAPHENE PER GALLON

ACTIVE INGREDIENTS:

*Toxaphene	70.6%
Aromatic Petroleum Distillate	24.5%

INERT INGREDIENTS:	4.9%
TOTAL	100.0%

E. P. A. Reg. No. 769-437

E. P. A. Est. 769-GA-1

*Technical Chlorinated Camphene (Chlorine content 67% to 69%)

WARNING! KEEP OUT OF REACH OF CHILDREN.

May be fatal if swallowed! Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after using. Do not store near feed or food products. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

DIRECTIONS

PROTECT WILDLIFE: This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PROTECT BEES: This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not feed treated crop or crop refuse to dairy animals or animals being finished for slaughter or to poultry. Do not allow grazing of treated crops by dairy animals or animals being finished for slaughter. Do not sell or ship forage treated with this material.

PECANS

To control pecan weevil, use at the rate of 1½ gallons to 500 gallons of water (24 pints per 100). Do not apply more than 300 gallons per acre. Follow locally issued spray schedule for number

and frequency of applications. Usually two applications are necessary, the first about July 30 in the latitude of Fort Valley, Ga., or when weevils begin to appear as determined by jarring, and a second application should follow in 10 days to 2 weeks. Do not apply after shucks begin to open. Do not graze livestock in treated groves.

LIVESTOCK

Do not use on dairy animals. Do not use in excess of recommended rate. Concentrations of twice the recommended rate may be fatal to young animals. Do not apply to animals within 28 days of slaughter. Do not use this product in dips. Do not use on calves under 3 months of age. Avoid treating animals in cold stormy weather, and overheated or sick animals.

BEEF CATTLE: Horn flies, Lice and Ticks - Use 4 pints per 100 gallons of water (6 oz. to 5 gallons of water) and spray at the rate of 2 quarts per animal. Apply at 3 week intervals or as needed for horn flies and lice.

SHEEP AND GOATS: Lice, Keds (Sheep tick) - Use 4 pints per 100 gallons of water (6 oz. to 5 gallons of water) and spray thoroughly. Repeat treatment in 2 to 3 weeks if necessary.

VEGETABLES

Do not apply to or allow drift to reach cucumbers, squash, or melons or injury may result.

COWPEAS: Cowpea curculio - Apply at the rate of 3 pints per acre in sufficient water for coverage. Make the first application at the time of bloom or when first small peas appear. Make 2 to 4 applications at 4 to 5 day intervals. Do not apply within 10 days of harvest. Apply only to peas that will be shelled. Do not apply to peas to be used as green snaps after pods begin to form. Do not feed treated vines to dairy animals or animals being finished for slaughter.

TOMATOES, PEPPERS, EGGPLANT: Hornworms, Blister beetles, and Pepper weevils on peppers - Apply at the rate of 3 pints per acre on peppers and eggplant and 2 to 4 pints on tomatoes in sufficient water for coverage. Do not apply to tomatoes within 7 days of harvest or to peppers and eggplant within 5 days of har-

vest. Remove all residue at harvest by washing or other effective means.

CUTWORMS: On beans, broccoli, cabbage, cauliflower, celery, eggplant, kale, peppers and tomatoes - Apply 2 pints per acre in sufficient water for coverage to soil surface before planting or to soil surface immediately after setting plants. Do not apply when plants are past seedling stage.

COTTON

THRIPS: For early season control, apply at the rate of ¾ to 1½ pints per acre in sufficient water for coverage, usually 2 to 3 gallons with one nozzle per row. Make the first application as soon as cotton is up to a stand and a second application 7 days later.

SOYBEANS AND PEANUTS

CUTWORMS, FALL ARMYWORM, VELVETBEAN CATERPILLAR: Apply at the rate of 2½ pints per acre in sufficient water for coverage. Make application as soon as small worms are found. Do not apply to soybeans within 21 days of harvest.

BARLEY, OATS, WHEAT, RYE, CORN, SORGHUM ARMYWORM, FALL ARMYWORM, and GRASSHOPPERS: Apply at the rate of 2½ pints per acre in sufficient water for coverage. Do not apply to sorghum within 28 days of harvest. Do not apply to sorghum more than once after heads begin to form. Do not ensile treated forage.

DISPOSAL WARNING

Dispose of waste pesticide and spillage by burying in non-crop land away from water supplies. Containers should be perforated or crushed and buried with waste.

NOTICE

DISCLAIMER OF WARRANTY, MERCHANTABILITY AND LIABILITY

If for any reason this product is not used, applied, stored, transported or disposed of in accordance with the instructions set forth on this label, then and thereafter Woolfolk Chemical Works, Inc. terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness.

WOOLFOLK CHEMICAL WORKS, INC.

MANUFACTURERS

470327

FORT VALLEY, GEORGIA