SECURITY

METHYL PARATHION EC-6

CONTAINS 6 POUNDS METHYL PARATHION PER GALLON

ACTIVE INGREDIENTS:

O, O-dimethyl O-p-nitrophenyl phosphorothioate ..... [illegible]

Aromatic Petroleum Distillate ..... [illegible]

INERT INGREDIENTS: ..... [illegible]

E. P. A. Reg. No. 769-440 TOTAL [illegible]

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME

POISON

ANTIDOTE AND FIRST AID TREATMENT

DANGER! KEEP OUT OF REACH OF CHILDREN. Poisonous if swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin. Repeated inhalation or skin contact may, without symptoms, progressively increase susceptibility to Methyl Parathion poisoning. Do not swallow or get in eyes, on skin or on clothing. Do not breathe mist or vapor. Wear clean natural rubber gloves, clean waterproof protective [illegible] and goggles. Replace gloves frequently and destroy used gloves. Wear a [illegible] respirator of a type passed by the U. S. Bureau of Mines for Methyl Parathion protection. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into the eyes, immediately flush with water for at least ten minutes and get [illegible] attention. In case of contact, remove all contaminated [illegible] wash skin with soap and water, wash clothing before re-[illegible]

PROTECT WILDLIFE. This product is toxic to fish, [illegible] other wildlife. Birds feeding on treated areas may be [illegible] of lakes, streams, or ponds. Do not apply where [illegible] cur. Do not apply when weather conditions favor drift [illegible] Shrimp and crab may be killed at application rates [illegible] label. Do not apply where these are important [illegible] taminate water by cleaning of equipment [illegible] this product only as specified on this label.

[illegible] This product is highly [illegible] direct treatment or residues on crops. [illegible] tained from your Cooperative Agricultural Extension [illegible]

ATROPINE IS THE EMERGENCY ANTIDOTE FOR METHYL PARATHION POISONING. [illegible] antidotal and may be administered [illegible] with atropine [illegible] your physician about obtaining an adequate supply of [illegible] gram atropine tablets for emergency use. CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING. **INTERNAL:** If the material has been swallowed, [illegible] immediately. This may be done by [illegible] with salt water [illegible] tablespoon of salt to a glass of water. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. **EXTERNAL:** If the material has been [illegible] remove patient from the [illegible] Methyl Parathion [illegible] and wash [illegible] water [illegible] the eyes, wash immediately with [illegible] water [illegible]

IF WARNING SYMPTOMS APPEAR: [illegible] gram atropine tablets immediately. NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR. [illegible] Keep patient prone and quiet. Start artificial respiration [illegible] patient [illegible] breathing. Transport the patient immediately to the nearest [illegible]

NOTE TO PHYSICIAN

WARNING SYMPTOMS: [illegible] weakness, headache, tightness [illegible] sweat [illegible]

TREATMENT: [illegible]

DIRECTIONS

[illegible]

COTTON: Boll Weevil, Aphids, Cotton Leafworm, Fleahopper, Thrips, and some species of Spider Mites - Apply at the rate of 1 2 to 2 3 pints per acre in sufficient water for coverage. Begin applications when insect infestation warrants and repeat at 4 to 5-day intervals as necessary for control.

BOLLWORM: This product will not control bollworms at the above rate. When this pest is present, increase the rate of Methyl Parathion EC-6 to [illegible] fluid ounces per acre. Do not apply this product within 7 days of hand picking or harvest. Cotton which has been treated with this material should not be grazed.

VEGETABLES: Aphids, Armyworms (up to 3rd instar), Flea Beetles, Leafhoppers and certain species of Mites. On Celery, Peas, Peppers, Potatoes, Lettuce, Spinach - Apply at the rate of 2 3 pints per acre. On Carrots, Beets, Tomatoes - Apply at the rate of 2 3 pints per acre. Do not use treated carrot tops for food or feed. On Beans, Cole Crops, Broccoli (Brussels Sprouts, Cabbage and Cauliflower) - Apply at the rate of [illegible] pints per acre. Begin applications when insects appear; repeat as necessary for control. Do not apply to Potatoes within 5 days of harvest, to Carrots, Celery, Peas, Peppers, Beets (where tops are not used for food), Beans (except green beans), and Tomatoes within [illegible] days of harvest, or to Beets, Lettuce, Spinach, Green Beans [illegible] Crops within [illegible] days of harvest.

ALFALFA: Spotted Alfalfa Aphids - Apply at the rate of 5 fluid ounces per acre in early spring. Aphids, Armyworms (up to 3rd instar), Leafhoppers and certain species of Mites - Apply at the rate of 2 3 pints per acre. Do not apply within 30 days of harvest.

SMALL GRAINS: Barley, Oats, Wheat - Aphids (Green Bug) - Apply at the rate of 5 to 8 fluid ounces per acre. Do not apply after heads begin to form.

DISPOSAL WARNING

[illegible] equipment and containers and dispose of wastes and spillage by [illegible] land away from water supplies. Never re-use containers [illegible] should be disposed of by punching holes in them and burying with wastes.

DISCLAIMER OF WARRANTY, MERCHANTABILITY AND LIABILITY

[illegible] this product is not used, stored, applied, transported or [illegible] accordance with the instructions set forth on this label, [illegible] thereafter Woolfolk Chemical Works, Inc. terminates, ends and [illegible] warranties, expressed or implied, including the warranty of merchantability and fitness.

371130

Manufacturers **WOOLFOLK CHEMICAL WORKS, INC.** **Ft. Valley, Ga.**