







2-(thiocyanomethylthio) INERT INGREDIENTS: .	benzothiazole		30% 70%
E.P.A. Reg. No. 769-446	E.P.A. Est. 769-GA-1	TOTAL	100%

## KEEP OUT OF REACH OF CHILDREN.

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID TREATMENT: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wesh contaminated clothing before re-use. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or by disposal of waste.

## **GENERAL INFORMATION**

WOOSAN 30 is recommended for the control of soil- and seed-borne fungi which impair the germination and seedling development of field corn, oats, drill-seeded rice, and wheat. WOOSAN 30, when used as directed, will usually increase stands and yields of field corn, oats, drill-seeded rice, and wheat by preventing seed decay and seedling diseases.

WOOSAN 30 can also be used for the control of covered smut of barley, Helminthosporium stripe of barley, loose and covered smut of oats, and stinking smut or bunt of wheat.

Do not use crops grown from WOOSAN 30 treated seed for grazing of livestock


WOOLFOLK CHEMICAL WORKS, INC. MANUFACTURERS 471112

or to hervest as silage or forage. However, grain harvested from plants grown from WOOSAN 30 treated seed can be used for food or feed. Label treated seed. Seeds treated with WOOSAN 30 cannot be used for food. feed, oil production, or any other purpose except planting.

FIELD CORN: Seed Protection - For ready-mix treaters, use 2.0 fluid ounces of WOOSAN 30 per 100 pounds of corn seed. WOOSAN 30 may be dispersed in water for distribution, provided the recommended amount of product is applied and the dispersion is agitated. For slurry treaters, disperse 2.0 fluid ounces of WOOSAN 30 in 10 fluid ounces of water and apply this dispersion per 100 pounds of seed. However, the concentration and amount of slurry to be used must be adjusted to the specific conditions of the slurry treater, provided that 2.0 fluid ounces of WOOSAN 30 is applied per 100 pounds of seed.

bushel of seed.

If for any reason this product is not used, applied, stored, transported or disposed of in accordance with the instructions set forth on this label, then and thereafter Woolfolk Chemical Works, Inc. terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness.



BARLEY, OATS, DRILL-SEEDED RICE, AND WHEAT: Seed Protection -For ready-mix treaters, use 0.75 fluid ounces of WOOSAN 30 per bushel of seed. WOOSAN 30 may be dispersed in water for distribution, provided the recommended amount of product is applied and the dispersion is agitated. For slurry treaters, disperse 0.75 fluid ounces of WOOSAN 30 in 4 fluid ounces of water and apply this dispersion per bushel of seed. However, the concentration and amount of slurry to be used must be adjusted to the specific conditions of the slurry treater, provided that 0.75 fluid ounces of WOOSAN 30 is applied per

Do not re-use container. Destroy when empty.

## DISCLAIMER OF WARRANTY, MERCHANTABILITY AND LIABILITY

## FLAT VALLEY, GEORGIA