

SECURITY

# 4% DIAZINON DUST

**ACTIVE INGREDIENT:**

O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate . . . . . 4.0%  
**INERT INGREDIENTS:** . . . . . 96.0%

E. P. A. Reg. No. 769-452      E. P. A. Est. 769-GA-1      TOTAL 100.0%  
Diazinon® is a trademark of CIBA-GEIGY Corporation.

**CAUTION!** KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. May be absorbed through skin. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Avoid breathing of dust.

**ENVIRONMENTAL CAUTION:** This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of waste. Apply this product only as specified on this label.

**PROTECT BEES:** This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

## DIRECTIONS

In the garden, apply Security 4% Diazinon® Dust when insects first appear and repeat applications as necessary for control. Dust for thorough coverage. Follow time intervals between last application and harvest. To control Mexican Bean Beetle, direct dust to underside of leaves.

### VEGETABLES

**BEANS (pole beans, snap beans, lima beans):** To control Black Bean Aphids, Mexican Bean Beetle, Leaf Miners, Mites, and Cucumber Beetles, apply 30 to 35 pounds per acre (1½ to 2 ounces of dust per 50 feet of row). Do not apply within 7 days of harvest.

**BROCCOLI, CABBAGE, and CAULIFLOWER:** To control Aphids, Diamondback Moths, and Imported Cabbageworm, apply 30 to 40 pounds per acre (1½ to 2 ounces of dust per 50 feet of row). Do not apply within 5 days of harvest to Broccoli or Cauliflower, or within 7 days of harvest to Cabbage.

**BRUSSELS SPROUTS:** To control Aphids, Diamondback Moths, and Imported Cabbage worms, apply 20 pounds per acre (1 ounce of dust per 50 feet of row). Do not apply within 7 days of harvest.

**COLLARDS, KALE, SWISS CHARD, and TURNIP TOPS:** To control Aphids, Imported Cabbageworms, Diamondback Moths, Harlequin Cabbage Bugs, and Corn Earworms, apply 30 to 40 pounds per acre (1½ to 2 ounces of dust per 50 feet of row). Do not apply within 10 days of harvest to Collards, Kale, or Turnip Tops, or within 12 days of harvest to Swiss Chard.

**CUCUMBERS:** To control Aphids, Serpentine Leaf Miners, and Thrips, apply 40 pounds per acre (2 ounces of dust per 50 feet of row). Do not apply within 7 days of harvest.

**MELONS and SQUASH (Summer and Winter):** To control Aphids, Leafhoppers, Mites, Ser-

pentine Leaf Miners, and Thrips, apply 30 to 35 pounds per acre (1½ to 2 ounces of dust per 50 feet of row). Do not apply within 3 days of harvest to Melons or Winter Squash, or within 7 days of harvest to Summer Squash.

**ONIONS:** To control Onion Thrips, apply 25 to 30 pounds per acre (1½ ounces of dust per 50 feet of row). Do not apply within 10 days of harvest.

**PEAS:** To control Aphids and Leaf Miners, apply 12½ pounds per acre (½ ounce of dust per 50 feet of row). May be applied up to day of harvest. Wait 4 days following treatment if pea vines are used for forage. Pea vines and pea vine hay may be fed to dairy cattle, beef cattle, and sheep.

**PEPPERS:** To control Aphids and Serpentine Leaf Miners, apply 25 to 40 pounds per acre (1½ to 2 ounces of dust per 50 feet of row). Do not apply within 5 days of harvest.

**POTATOES:** To control Aphids, Flea Beetles, Colorado Potato Beetles, and Dipterous Leaf Miners, apply 30 to 40 pounds per acre (1½ to 2 ounces of dust per 50 feet of row). To control Banded Cucumber Beetles, Leafhoppers, and Southern Armyworms, apply 40 to 50 pounds per acre (2 to 2½ ounces of dust per 50 feet of row). Do not apply within 35 days of harvest.

**SPINACH and BEETS:** To control Aphids and Dipterous Leaf Miners, apply 20 pounds per acre (1 ounce of dust per 50 feet of row). Do not apply to Spinach within 10 days of harvest or to Beets within 14 days of harvest.

**SWEET CORN:** To control Corn Earworms, apply when silks first appear. Two to three applications are usually necessary at 2 to 3 day intervals. Apply 30 to 40 pounds of dust per acre (1½ to 2 ounces of dust per 50 feet of row). Do not apply within 2 days of harvest. Do not feed treated forage to animals being finished for slaughter within 2 days after treatment.

**TOMATOES:** To control Aphids, Dipterous Leaf Miners, Banded Cucumber Beetles, Fall Armyworms, and Southern Armyworms, apply 30 to 40 pounds per acre (1½ to 2 ounces of dust per 50 feet of row). Do not apply within 3 days of harvest.

### FRUITS

**BLUEBERRIES:** To control Blueberry Maggots, apply 25 pounds per acre. Do not apply within 7 days of harvest.

**GRAPES:** To control Grape Berry Moths, Grape Leaf Folders, and Leafhoppers, apply 20 to 25 pounds per acre. Do not apply within 10 days of harvest.

**STRAWBERRIES:** To control Aphids, Two-Spotted Mites, and Strawberry Leaf Roller, apply 20 to 30 pounds per acre. To control Cyclamen Mites, apply 30 pounds per acre and direct dust to plant crowns. Do not apply within 5 days of harvest.

### DISPOSAL WARNING

Do not re use container. Destroy it by burying or burning. Avoid smoke or fumes.

### NOTICE - Disclaimer of Warranty, Merchantability, and Liability

If for any reason this product is not used, applied, stored, transported or disposed of in accordance with the instructions set forth on this label, then and thereafter Woolfolk Chemical Works, Inc. terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness. 570507

Manufacturers **WOOLFOLK CHEMICAL WORKS, INC.** Ft. Valley, Ga.