SECURITY ACCEPTED

4% Diazinon Dust

NET WEIGHT 25 POUNDS

ACTIVE INGREDIENT

O,O-Diethy O [illegible] phosphorothioate [illegible]

INERT INGREDIENTS [illegible]

EPA Reg. No. 769-4[illegible] EPA Est. 769-GA-[illegible]

Diazinon® is a registered trademark of CIBA [illegible] Corporation

KEEP OUT OF REACH OF CHILDREN.

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED Drink one to two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Call a physician immediately.

IF INHALED Remove victim to fresh air and apply respiration if needed.

IF ON SKIN: Wash with soap and water.

IF IN EYES Flush eyes with water for 15 minutes and call a physician immediately.

NOTE TO PHYSICIAN If signs of cholinesterase inhibition are present atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine sulphate.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

[illegible]

ENVIRONMENTAL HAZARDS

[illegible]

PROTECT BEES [illegible]

DIRECTIONS FOR USE

[illegible] with its [illegible]

[illegible] the garden apply [illegible] pear and repeat applications [illegible] mechanical or hand duster [illegible] parts. Dust for thorough [illegible] plication and harvest [illegible] derside of leaves.

VEGETABLES

Beans (pole beans, snap beans, lima beans) [illegible] Mexican bean beetle [illegible] [illegible] pounds of dust per [illegible] days of harvest.

Beets [illegible] within 7 days of harvest.

Broccoli, Cabbage and Cauliflower To control aphids [illegible] cabbageworm apply 30 to 40 pounds [illegible] feet of row. Do not apply within [illegible] days of harvest.

Brussels Sprouts [illegible] cabbageworm apply [illegible] days of harvest.

Collards, Kale, Swiss Chard, and Turnip Tops To control aphids, imported cabbageworm, diamondback moths, Harlequin cabbage bugs and corn earworms, apply 30 to 40 pounds [illegible] 10 2 ounces of dust per 50 feet of row. Do not apply within 10 days of harvest to collards, kale, or turnip tops or within 12 days of harvest to swiss chard.

Cucumbers To control aphids, serpentine leaf miners and thrips, apply 40 pounds per acre (2 ounces per 50 feet of row). Do not apply within 7 days of harvest.

Melons and Squash (Summer and Winter) To control aphids, leafhoppers, mites, serpentine leaf miners and thrips, apply 30 to 35 pounds per acre (1 to 2 ounces per 50 feet of row). Do not apply within 3 days of harvest to melons or winter squash or within 7 days of harvest to summer squash.

Onions To control [illegible] feet of row. Do not apply within [illegible] days of harvest.

Peas [illegible]

Peppers [illegible]

Potatoes [illegible]

Spinach and Beets [illegible]

Sweet Corn [illegible]

Tomato [illegible]

SOIL APPLICATION

[illegible] Potatoes
[illegible] Melons, Squash, Spinach
[illegible] Tomatoes
[illegible] Parsnips

Apply as a broadcast treatment over the surface of the soil. Apply at the rate of 1 pound per 450 square feet. Immediately after spreading, mix thoroughly with the soil to a depth of 6 to 8 inches.

To control ROOT MAGGOT in onions, apply in furrow at the rate of 9 tablespoons per 100 feet of row at the time of planting.

To control ROOT MAGGOT in radishes, apply in furrow at the rate of 9 tablespoons per 100 feet of row at the time of planting.

SMALL FRUITS

BLUEBERRIES To control blueberry maggots, apply 25 pounds per acre. Do not apply within 7 days of harvest.

GRAPES To control grape berry moths, grape leaf folders and leafhoppers, apply 20 to 25 pounds per acre. Do not apply within 10 days of harvest.

STRAWBERRIES To control aphids, two-spotted mites and strawberry leaf roller, apply 20 to 30 pounds per acre. To control cyclamen mites, apply 30 pounds per acre and direct dust to plant crowns. Do not apply within 5 days of harvest.

ROSES, FLOWERS, AND SHRUBS

To control aphids, bagworms, mites, flea beetles, leafhoppers, leafminers, mealybugs, webworms, whiteflies, tent caterpillars, scale crawlers and [illegible] repeat as necessary. For best [illegible] when the temperature is low [illegible] parts of plants are contacted [illegible] use on terns [illegible] not effective [illegible]

LAWN INSECTS

Spot Treatment [illegible] wasps [illegible] thoroughly on and around [illegible] when wasps are not active.

Band Treatment Around Foundation [illegible]

DISPOSAL

NOTICE [illegible] the Warranty [illegible]

12

Voolfolk Chemical Works, Inc.
Manufacturers Ft. Valley, Ga.