



PurgeTM Weed Killer

FOR ST. AUGUSTINE GRASS
CENTIPEDE GRASS
AND DORMANT BERMUDA GRASS

ACTIVE INGREDIENTS

Atrazine 10.0%
Alachlor 10.0%
Glyphosate 10.0%

INERT INGREDIENTS

Water 70.0%
Surfactant 10.0%
This product contains
Attention: Do not spray

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

ACCEPTED

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION

REGIONS FOR USE

GENERAL INFORMATION

APPLICATION

KEEPING AND STORAGE

HOUSEHOLD APPLICATION

NET CONTENTS

FLORIDA BETONY This weed emerges in the fall. An application of Purge in mid to late October followed with a second application in late February would give the best control.

DICHONORA MONETWORT The best control for these weeds can be obtained by applying Purge in early April followed with a second application in late May. In growing bermuda grass areas, bermuda grass will generally give control of said weeds.

FLORIDA BETONY This weed emerges in the fall. An application of Purge in mid to late October followed with a second application in late February would give the best control.

ST. AUGUSTINE AND CENTIPEDE GRASSES

Apply Purge to St. Augustine and Centipede grasses in late August or early September. Best results are obtained when weeds are small and the grass is in the growing season.

DORMANT BERMUDA GRASS

Apply Purge to dormant bermuda grass during the dormant season. Best results are obtained during the growing season. Cause in

DATE OF APPLICATION

Apply Purge in mid to late October followed with a second application in late February. State contents vary.

LISTED AMOUNT

1 gallon
2 gallons
3 gallons
4 gallons
5 gallons

DISPOSAL

Do not dump in waterways or on land. Dispose of in accordance with local regulations.

© 1987, Chemagro Corp., Norfolk, Virginia